

A Survey of Pertinent Biochemical Literature

By EUGENE T. OBORN

CHEMISTRY OF IRON IN NATURAL WATER

GEOLOGICAL SURVEY WATER-SUPPLY PAPER 1459-F



UNITED STATES DEPARTMENT OF THE INTERIOR

FRED A. SEATON, *Secretary*

GEOLOGICAL SURVEY

Thomas B. Nolan, *Director*

CONTENTS

	Page
Abstract.....	111
Introduction.....	111
Nontechnical definition of terms.....	113
Biochemical factors in the iron cycle.....	114
Biochemical weathering.....	114
Iron metabolism.....	115
Relation of iron to life processes.....	117
Solution and deposition of iron.....	118
Organisms utilizing iron in metabolism.....	124
Fungi.....	124
Bacteria.....	124
Protozoa.....	125
Phages.....	125
Viruses.....	125
Yeasts.....	125
Algae.....	126
Micro-organisms, general.....	126
Flowering plants.....	127
Animals.....	128
Organisms, general.....	128
Iron bacteria.....	129
Mechanics of iron concentration and depletion.....	134
Lichen effects.....	134
Hydrophyte (other than bacteria and fungi) effects.....	134
Chemical elements interfering with use of iron by plants.....	141
Measurement of materials affecting iron solubility.....	143
Analytical ramifications of iron biochemistry.....	143
Selected references.....	149
Index.....	189

ILLUSTRATIONS

	Page
FIGURE 12. Tearing away of glass surface by drying contraction of 2- and 5-percent solutions of the plant gelatin agar-agar	116
13. Diagrams of different iron bacteria	130
14. Relative importance of the gametophyte and sporophyte generations in the four major groups of aquatic plants	135
15. Morphological comparison between leaf-midrib and root structures of rice and those of typical aquatic and land plants	136
16. Some phases of the iron cycle that affect the presence or absence of iron in natural water	139
17. Regions of dicotyledon and monocotyledon plants in which iron and other metallic salts tend to concentrate	140

TABLES

	Page
TABLE 1. Organisms that require iron for normal metabolism	133
2. Humic matter, dissolved oxygen, and tannic analyses	144
3. Methods of analysis for iron in water, soil, and plant material ...	145

CHEMISTRY OF IRON IN NATURAL WATER

A SURVEY OF PERTINENT BIOCHEMICAL LITERATURE

By EUGENE T. OBORN

ABSTRACT

Biochemical factors relating to the occurrence of iron in natural water were studied by examining more than 800 published articles and abstracts. According to this literature iron is essential in normal metabolism of most of the fungi, bacteria, aquatic vegetation, and land plants studied as well as in many animals. Many organisms in diverse habitats may precipitate or dissolve iron. Aquatic plants seem to be important in the iron cycle in surface-water bodies. They take up iron from submerged soil or from the water and ultimately release it to water solution or return it to the bottom mud through rapid decay of dead plant material. Anaerobic bacteria in the absence of oxygen reduce precipitated ferric iron and return it to solution. Apparently, many of these bacteria in the presence of oxygen use the carbon that is so frequently fixed to iron in the form of humates, chelates or other complex compounds. Crustose lichens remove iron from bare rocks. Other life forms are also involved in the iron cycle.

Under oxidizing and alkaline conditions, mud layers in lake bottoms act as an adsorbent for the more common insoluble metallic ions. Mud layers reduced by anaerobic bacteria release iron and the more common metallic ions. The simpler aquatic plants use these metallic ions directly.

The redox potential of water is related to the content of dissolved ferrous and ferric iron. Iron in organic complexes may be difficult to determine by usual methods of chemical analysis. Special methods and techniques are suggested when the determination is to be performed in the presence of organic matter.

INTRODUCTION

This survey of published literature was made to aid further study and to summarize the knowledge of chemical and (or) biochemical factors that govern the solution, transport, and deposition of iron by natural water. No comprehensive summary of the subject has been published. However, many scientists have studied certain phases of the iron cycle in nature. Much of this work concerns the use of iron by plants and animals. This iron use is directly related to the occurrence of iron in water.

Iron is undesirable in water for human use. In amounts exceeding a few tenths of a part per million, iron in domestic water may stain porcelain fixtures, glassware, and laundered articles and may impart an unpleasant taste to water or to foods prepared with such water. In water for industrial use, iron may interfere with textile bleaching, dyeing or finishing, paper production, and manufacture of wines, beer, and other beverages; it may also impair the quality of dairy products or interfere with fish propagation. Therefore iron removal is essential in water treatment (Nordell, 1953).

Chemical-element cycles, particularly those of nitrogen and carbon, are presented in many elementary plant-science textbooks. These cycles emphasize the changes in elemental combination caused by microscopic and macroscopic organisms. Some phases of the iron cycle that affect iron content in natural water can be simply explained in diagrams.

More than 800 articles and abstracts were examined. These are listed in the bibliography, and many are summarized in this report. Common names used in the original articles are cited along with scientific names unless only one name was available.

In order to review thoroughly and systematically the factors affecting iron content in natural water, it was necessary to investigate iron utilization by all kinds of life, particularly soil and aquatic flora. Because other ions (calcium, copper, and manganese to name just a few) affect iron utilization or solubility, the literature on foreign ion interference to iron solution and availability was pertinent. Utilization of iron by many soil organisms affirms the importance of organisms in removing iron from the soil. Organic material in water interferes with the accuracy of certain methods of iron analysis. Accordingly, published articles on methods of determining iron were examined. No attempt has been made to evaluate or classify the articles into basic or applied research.

Three works (Needham and Needham, 1935; Fassett, 1940; and Muenscher, 1944) are recommended for picture identification of the hydrophytes.

The interrelationship of water-soluble iron with other ions, pH, conductivity, redox potential, and dissolved oxygen is noted in many works and should be studied in any investigation concerning the solution of iron in and the removal of it from natural water.

The relationship of iron to chlorophyll, protein, ash, crude fiber, oxygen and carbon-dioxide production by both water-rooted and soil-rooted plants is important. Soil microflora leach iron from soils and affect the amount in natural water. The literature emphasizes the desirability of correlating and interpreting laboratory (greenhouse) findings with results obtained in the field (on a suitable lake).

NONTECHNICAL DEFINITION OF TERMS

The following selected biological terms are defined to facilitate the reading of this study:

Actinomycetes. Thread bacteria or fungi frequently breaking up into short rods.

Benthos. Organisms that inhabit the bottom of a lake or other water body.

Cambium. Perpetually young undifferentiated plant cells in perennial plants responsible for increased stem and root diameter giving rise to new tissue.

Chloroplast. An oval-shaped matrix in plant cells on which chlorophyll manufactured in the same plant cell is adsorbed.

Chromosome. Contains bent-rod appearing bodies made up of chromatin and probably present in all living cells.

Coenocyte. Organisms having cells with more than one nucleus.

Dicotyledon. Type of plant which may be identified in the field by presence of netted-veined leaves.

Gametophyte (haploid). Plant made up of cells having a single set of chromosomes. Conspicuous part of plant body as in many unicellular or filamentous algae, lichens and mosses.

Hydrophyte. Water plant.

Littoral. Inhabited water adjacent to the shore.

Meristem. Perpetually young undifferentiated plant cells responsible for differentiation of tissues in shoot, root, and leaf bases of plants.

Mesophyll. The green uniformly thin-walled plant cells between the epidermal layers of a foliage leaf.

Monocotyledon. Type of plant which may be identified in the field by presence of parallel-veined leaves.

Mycelium. Threadlike protoplasm containing filaments characteristic of fungi.

Nonvascular plant. Plant lacking a vessel or vessels for conveyance of sap through the plant. Mainly nonflowering plants such as the algae, fungi, lichen, bacteria, and moss groups.

Pelagic. Inhabiting open water.

Phage. Bacteria-destroying virus or agent normally present in sewage, the intestinal tracts of man and animals, and in blood, pus, and urine.

Phloem. Food-conducting tissue of flowering plants.

Plankton. Forms of life, frequently microscopic, near surface of open water.

Protozoa. Mostly microscopic one-celled animals.

Sporophyte (diploid). Plant made up of cells having a double set of chromosomes. Conspicuous part of plant body as in flowering plants and ferns.

Vascular plant. Plant having a vessel or vessels for conveyance of sap through the plant, mainly these are flowering plants, but include the ferns.

Virus. Submicroscopic infective substance capable of causing disease.

Xylem. Mineral-conducting tissue of flowering plants.

BIOCHEMICAL FACTORS IN THE IRON CYCLE

BIOCHEMICAL WEATHERING

Literature on the chemistry of iron in natural water emphasizes the widespread and continuous influence of biota on the amount and state of oxidation of iron in either moving or quiet water. All categories of fauna and flora, ranging in size from smaller than that seen through the regular biological microscope to tree-size growth, have many representatives which are vital in critical phases of the iron cycle.

Trees, shrubs, and garden crops are effective in removing iron from well-aerated soils. Fungi and particularly bacteria may remove or release iron from water-submersed or nonaerated soils. Rooted vascular hydrophytes are also prominent in the iron cycle and affect the solution of iron in natural water. The roots of this group of water plants generally grow in environments lacking oxygen. The plants are both emergent and submersed. They take oxygen to their own roots through special air tubes, and the roots absorb the relatively large amount of iron needed from the submerged mud. Unattached vascular hydrophytes, such as the surface floating duckweeds, are important in the iron cycle in natural water, particularly in ponds and in places protected from rapid waterflow, by (a) occluding normal air-water interchange of oxygen and (b) depleting the dissolved-oxygen supply of the water through respiration of rootlets projecting into and just below the water surface. Iron taken into solution in the resulting less-oxidizing medium is available for the needs of this plant. Mineral salts taken from the substratum are returned as organic mineral deposits upon death and decay of the aquatic organism (Butcher, 1933). Algae attached to bottom muds and sediments obtain most of the iron needed for metabolism directly from iron in the water in which the plants are submersed. The holdfast cells of fresh-water algae probably absorb some iron from the substratum to which they are attached; but in varieties lacking effective conduction tissue, most of the iron is absorbed by the individual generally haploid thallus cells of the filament as needed.

Some attached algae, such as the fresh-water stoneworts and the marine seaweeds, have functional conductive systems. These algae apparently absorb iron principally from the substratum rather than directly from the water.

Unattached or plankton algae obtain all the iron needed for metabolism directly from the water in which they are submerged. Apparently little or no iron is obtained directly from the substratum.

Crustose lichens are very active and effective in removing iron from barren rocks by physical and chemical methods. Further solution of iron from rocks is accomplished by successive lichen—foliose and fruticose—and moss growth. Fry (1924) believed that the rock substratum first was altered chemically by lichens and then the parts of the decomposed material were separated by the mechanical action of the growing hyphae. However, Fry has shown that these processes can be reversed, the initial stage being the mechanical disintegration of the unaltered rock by the lichen thalli, followed by the decomposition of the separated fragments. This mechanical action is illustrated in figure 12, which shows the tearing away of the surface of a piece of plate glass and of a petri dish by the drying contraction of 2- and 5-percent solutions of the plant gelatin agar-agar. Crustose and foliose lichens growing on schist are also illustrated. Lichens probably break up rock surfaces in a similar manner, one of the first steps in removing iron from the rock substratum to which they are attached. Gelatinizing agents, such as *d*-galactopyranose and *d*-galactose, through expansion-contraction wetting action bring about this mechanical disintegration of the rock substrata. Chemical disintegration comes about through carbonic and alginic acid solution and organic chelation processes (Schatz and others, 1956). It is noteworthy that the addition of gelatin to ferric solutions prevents (Taboury and Mangin, 1948) the precipitation of hydroxide when alkali is added.

IRON METABOLISM

Many have shown that iron in soluble or available form is indispensable for the synthesis of the chlorophyll-protein-lipoid complex in all green plants (Abadia, 1956; Samish, 1954; Singh, 1956). Organic-bound iron present as respiratory pigment in apparently all living protoplasm seems to act as a catalyst or oxygen carrier in oxidation-reduction processes occurring in living cells. Iron salts in any appreciable concentration, as a result of extreme soil acidity or great lack of aeration or both, are toxic to plants.

In spite of the absolute need of green plants for iron, the proportionate amount in plant tissues is very low and much of the iron pres-




FIGURE 12.—Tearing away of glass surface by drying contraction of 2-percent solution (in petri dish, lower left) and 5-percent solution (on plate glass, lower right) of the plant gelatin agar-agar. In similar manner crustose lichens (upper left) and foliose lichens (upper right) are believed to break up rock surfaces.

ent is a constituent of organic compounds. Most of the literature indicates that iron is immobile within plants; no appreciable redistribution ever occurs from one tissue to another after the tissues have been built up. Therefore, during growth, aquatic as well as land plants must have a continuous source of iron; iron extraction from the soil is a continuous process throughout the growing season. Aquatic plants average 90 to 95 percent water. Because many have no woody parts, disintegration at the end of the growing season is rapid. The resulting gelatinous byproducts, tannins, and other plant acids released tend to reduce and hold iron in solution until used by bacteria or other aquatic organisms. Death and disintegration of aquatic fauna that feed on aquatic flora likewise bring organic-bound reduced iron into solution.

RELATION OF IRON TO LIFE PROCESSES


Several fundamental life processes seem to depend on an adequate supply of iron (Ellis, 1919; Guelbenzu, 1951; Guseva, 1939; Kaserer, 1910; Rankama and Sahama, 1952; Steward, 1948; Warburg, 1925). Iron is needed for plant photosynthesis (which involves alternate oxidation and reduction of iron), respiration, and protein synthesis. In the higher animals, the mammals, iron forms an integral part of the hemoglobin or red-blood molecule. Amounts of iron in plants and animals are nearly parallel with amounts of chlorophyll or protein. Relationships of protein and chlorophyll synthesis (and amount of iron) are discussed by Bennett, 1945; Bonner, 1950; De Boissezon, 1933; Escudero and others, 1944; Kliman, 1937; Peterson and Elvehjem, 1928; Randoin and Le Gallic, 1942; Sherman, 1907; Sisakyan and Melik-Sarkisyan, 1956; Williams, 1953; Wolken and Mellon, 1956.

Loss of iron from solution by precipitation as hydroxides and the effect of complexing in organic compounds normally limit the dissociated iron content of natural water to levels within tolerance limits for plant growth (Smith, 1950). If iron concentrations in water are excessive or when plant tissue comes directly in contact with certain iron compounds, as for example iron sulfate sprayed on young plants when they have their first 2 or 3 leaves, vegetation is destroyed by corrosive action (Duchaufour, 1953a). Before the advent of growth regulators, this principle was commonly applied in weed control (Adams, 1909; Beaumont and Holland, 1935; Blum, 1956; Blunno, 1911; Clark, N. A. 1936; Coste, 1896; De Jesus, 1937; Dusserre, 1900; Graftiau, 1900; Handke, 1933; Kharasch and others, 1936; Maze and Perrier, 1904; Ponnampereuma and others, 1955; Robinson 1930, 1951; Setter, 1930; Smith, 1950; Voicehovich, 1940; Willis, 1936; Yoshimura, 1936).

Most aquatic plants grow actively in the young tissue by rhizomes and die in the old tissue; therefore, they remain perpetually young (Arber, 1920). Thus, particularly in these plants, an extra large amount of perpetually young undifferentiated tissue requires iron for normal metabolism.

Because of the difference in methods of iron absorption, most but not all accumulation or concentration is accomplished by the lower algal forms of the hydrophytes. This process withdraws some of the dissolved matter of the water body.



Although plant life alters the iron content of natural water, animal life also alters the content through direct or indirect dependence on plants. More directly, mankind probably derives much of its daily iron need from culinary water (Asbury in Forman and Frink, 1950; Sebrell, written communication, 1942). The empirical chemical formulas of photosynthesis and respiration are as follows: Photosynthesis (restricted to cells containing chlorophyll).




Aerobic respiration (living cells breathing atmospheric oxygen).




Anaerobic respiration (living cells not breathing atmospheric oxygen).




or



or



Molecular hydrogen is released by the hydrogenase of the mud organism and (or) organisms as in the reaction



Apparently, most of the carbon dioxide used in photosynthesis may be extracted from the soil (Dyakonova, 1957), and most of the carbon dioxide of underground carbonic acid water may originate in the atmosphere (Smirnov, 1955).

SOLUTION AND DEPOSITION OF IRON

Reducing and acid conditions (Goldschmidt, 1954), especially if pH is less than 3, generally promote iron solution in water. With the semiannual decrease of dissolved oxygen in lake bottoms, which in new reservoirs or in freshly inundated marshland may amount to total depletion, activity of anaerobic bacteria and other organisms

results in the solution of iron and manganese from the bottom deposits. Organic color in water represents the extract of soluble material contained in decaying vegetation, such as leaves and debris in peat bogs. Generally, the color is a brownish yellow, becoming deeper brown at higher concentrations. Iron may be an integral constituent in this organic color in water; and upon oxidation of the organic matter, the iron is precipitated and settles to the bottom of the body of water. If, during summer stagnation, conditions become anaerobic, the iron may be redissolved and later distributed throughout the water as colloidal ferric hydroxide. This change, however, is not a permanent condition; sedimentation and clarification soon follow.

The following biochemical processes or combination of processes aid solution of iron in natural water:

1. Formation (Rummeni, 1955) of citrate, tartrate, malate, and humate complexes of ferric iron (citric, malic, lactic, and tartaric acids, hemoglobin, chlorophyll, urine, and manure are chelating agents).
2. Reduction of iron to the ferrous state.
3. Processes releasing carbon dioxide or (other) sources of hydrogen ions.
4. Formation of complex ions, such as those with sodium hexametaphosphate.
5. Possible prevention by organic matter of formation of iron nuclei needed for crystallization.

Water draining from mines may be acidic, and being dependent on the state of oxidation of the iron (Braley, 1957), may contain large amounts of ferrous and ferric iron owing to oxidation of pyrite. Pyrite (FeS_2) deposited under conditions of oxygen deficiency (Murray, 1956), is oxidized (Segeberg, 1937) to ferric oxide and sulfuric acid in the presence of air and water.

Carbonic acid resulting from both aerobic and anaerobic respiration is an important aid in solution of iron in soil moisture and thus makes iron available, frequently as the bicarbonate, for absorption by rooted green plants (Shilova and Kreier, 1957). The preceding is true even though in well-drained and well-aerated soils oxygen is available in the plant root zone.

Many publications show organic matter to be an aid to iron solubility. Processes of altering iron solubility with the aid of organic matter are discussed by: Aiyar, 1946, 1948; Aleev and Mudretsova, 1937; Alexander and Walsh, 1952; Allyn, 1927; Betremieux, 1951; Beutelspacher, 1955; Bhaskaran and Pillai, 1937; Biddulph, 1948; Bloomfield, 1952, 1957; Boiret and Paturel, 1892; Boischot and Dur-

roux, 1949; Bould, 1957; Brewer and Carr, 1927; British Standards Institute, 1956; Broadbent, 1954; Carnes, 1954; Chaberek and Bersworth, 1953; Clark and others, 1957; Clark, N. A., 1936; Coldwell and De Long, 1950; Cooper, 1948; Correns, 1941; Datsko, 1955; De Kock, 1955a, 1955b; De Long and Schnitzer, 1955; Demolon and Bastisse, 1938; Doyne and Morison, 1926; Duchaufour, 1953a; Edelstein and Von Csonka, 1912; Eggleton, 1931; Falci, 1935; Fly, 1935; Gellerman, 1950; Gile and Carrero, 1916b; Guelbenzu, 1951; Hance, 1934; Hopkins and Wann, 1925, 1926; Hutchinson, 1941; Ichikawa, 1936; Iyengar, 1936; Iyengar and Subrahmanyam, 1935; Joffe and McLean, 1928; Kahler, 1956; Kapp, 1932a, 1932b, 1934a, 1934b; Kaserer, 1910; Kawamura, 1928; Kliman, 1937; Koeck, 1956; Kopal, 1940; Kuhnholz-Lordat, 1928; Lackey, 1939; Leonard and Stewart, 1953; Lipmann, 1937; Loew, 1938; Lutwick and De Long, 1954; Lutwick and others, 1952; Malyuga, 1945; Mangenot, 1953; Martell and Calvin, 1952; Masoni, 1914; Mayer and Gorham, 1951; Maze and Perrier, 1904; Morison and Doyne, 1914; Nakabayashi, 1954; Narasinhmurthy, 1940; Narasinhmurthy and Subrahmanyam, 1953; Nierenstein, 1945; Ohle, 1937; Olivier and Le Peintre, 1955; Olsen, 1935; Onodera, 1928; Osugi and Aoki, 1936, 1938b; Osugi, Aoki and Morita, 1935; Osugi, Nishigaki, and Yoshimi, 1935; Pearsall, 1952; Pringsheim, 1949; Prozorovskaya, 1936; Ramaut, 1955; Rankama and Sahama, 1952; Reed and Haas, 1924; Robinson, 1930; Roelofs, 1944; Saint, 1935; Sarishvili and Bagaturiya, 1938; Saunders, 1938; Saunders, 1957; Schnitzer and De Long, 1954b, 1955; Schoeller and Webb, 1936; Seeborg, 1937; Segeberg, 1937; Siddappa and Subrahmanyam, 1935; Skeen, 1930; Skopintsev and Krylova, 1955; Smith, 1950; Steel and Gloyna, 1954; Stewart and Leonard, 1956; Tsujimura and Ikeda, 1957; Van Beneden, 1956b; Vialard-Goudon and Richard, 1955; Villeret, 1954a, 1954b; Vinberg, 1955; Wallace and Ashcroft, 1956; Wallace and others, 1955; Wallace, 1929; Wallace and Lunt, 1956; Weinstein and others, 1954; Weston, 1936; Whipple and others, 1927; Wiebosch, 1947; Willis, 1936; Yoshimura, 1931a.

Solubility of iron in soil affected by several interrelated biochemical phenomena, such as pH, adsorption, presence of calcium or sulfur, particle size, and different treatments of soil, is discussed by: Asana, 1946; Askinazi and Kheifets, 1938; Baker and Price, 1946; Bandt, 1938; Barbier, 1938a, 1938b; Barbier and Chabannes, 1949; Boisshot and Durroux, 1949; Brown and Holmes, 1955; Brozek, 1949; Bruevich and Vinogradova, 1946; Byram, 1938; Calgon and Albright & Wilson, 1944; California State Water Pollution Board, 1952; Clarke, 1924; Cooper, 1935; Delecourt, 1947; De Wilkoszewski, 1917; Dienert, 1938; Dion, 1944; Edgerton, 1942; Fischer, 1950; Follett-Smith, 1938;

Follett-Smith and Robinson, 1936; Goldschmidt, 1954; Haase, 1940; Hance, 1934; Harvard, 1937; Henry, 1950; Hesse, 1956; Hoore, 1949; Hseung, 1941; Hylkema, 1939; Ignatieff, 1939, 1941; Ingleson and Thomas, 1946; Joffe, 1930; Jolivet, 1934; Kapp, 1932a, 1932b; Keller, 1945; Keller, 1955; Klut, 1937; Koczy, 1950; Kriventsov, 1954; Kurzweil and Exner, 1954; Lackey, 1939; Levin, 1947; Luhrig, 1927; Lutz, 1936; MacCarthy, 1925; Mackenzie, 1952; Mandl, 1949; Mason, 1952; Metzger, 1930; Mohler, 1951; Mukherjee and Chatterjee, 1945; Mukherjee and others, 1946, 1947; Mulwani and Pollard, 1939; Navelier, 1950; Navet, 1947; Nordell, 1953; Notarov, 1956; Ohle, 1937; Okura and Kayama, 1952; Osugi and Aoki, 1938a, 1938b; Osugi and others, 1934; Oudin, 1938; Parker, 1955; Patrick, 1953; Pearsall, 1950; Pronin, 1933; Puri and Sarup, 1938; Puustjarvi, 1952; Rakestraw and others, 1936; Rippel, 1941; Ritter and Christen, 1935; Rudge, 1936; Saruhashi, 1955; Schachtschabel, 1942; Schilling, 1936b; Solovev, 1940; Speller, 1922, 1951; Thompson, 1940; Todt, 1934; Toth, 1937; Urs and others, 1955; Valyashko, 1955; Van Beneden, 1947, 1948; Walker and others, 1907, 1908; Wheatly, 1946; Whipple and others, 1927; White, 1937; Winters, 1940; Zonn, 1945.

In submerged soils lacking organic matter, reduction of iron or increases in cation concentration are not likely. Chemosynthesis taking place under ice (Sorokin, 1955) and organic matter resulting from the death and decomposition of the aquatic plants (Arber, 1920) do affect the redox potential of soils and natural water.

Mortimer (1942) and others have shown relationships between increasing depth and the fall in redox potential and oxygen concentration in lake water and the rise of color, iron, and turbidity. Increase in turbidity and color resulted mainly from ferric hydroxide in suspension. Oxidation of ferrous iron and other processes using up oxygen accounted for an accelerated decrease in oxygen concentration with depth. At the point of zero oxygen content ferrous iron was detected in the water, and the redox potential had fallen to 0.15 corrected to a pH of 7. From this point to the lake bottom, the iron content of the water increased rapidly, and an increasing proportion consisted of ferrous iron. When seasonal changes in temperature brought this bottom water to the surface, the water absorbed oxygen from the air, and then the iron concentration decreased rapidly because of the precipitation of ferric hydroxide.

When the oxygen concentration and redox potential of the water in contact with the bottom mud dropped below a limiting value, ferric iron at the mud surface was reduced to the more readily soluble ferrous form. Freshly precipitated ferric hydroxide has some capacity for adsorption of cations. The oxidized layer at the mud surface with its

adsorbed ions constitutes a barrier to exchange between the water above and below the layer.

Mortimer (1941) cites the work of Ohle (1937), who demonstrated the ampholytic nature of the ferric hydroxide gel, which is electropositive in acid solution and electronegative in alkaline solution. Thus, in the presence of carbon dioxide at pH 4, negatively charged phosphate ions added to ferric hydroxide gel were strongly adsorbed, although some of the iron went into solution. After raising the pH to 9, a large part of the phosphate was liberated. These phenomena partly explain the fundamental and somewhat paradoxical influence of the presence or absence of bases (calcium) on the biochemical activity in fresh water. In the water of calcium-poor humus in lakes, pH generally will be below 7 and negatively charged phosphate and other plant nutrients are adsorbed on positively charged iron gels, which are themselves attached to electronegative humus colloids and are retained in the bottom muds; hence, few organisms can be supported.

An observation made by Mortimer (1941) and supported by many laboratory studies is that ferrous iron in water under natural conditions is not oxidized instantaneously. This fact was shown by conditions found during mixing preceding the overturn of lake water. At this time, ferrous iron was detected in water containing 8.4 milligrams per liter of oxygen. The rapid spread of ferrous ions in the water above the mud layer was believed by Mortimer to be the result of transition from molecular diffusion in the mud to turbulent diffusion in the water. The ferrous ions were oxidized and precipitated; thus the oxygen was depleted and an oxidized layer was built up at the surface of the bottom mud. When anaerobic conditions were re-established at the bottom, the adsorbent oxidized surface mud layer became progressively thinner and disappeared about 40 days after oxygen was depleted. A rise in temperature accelerates greatly the rate of decomposition of organic matter. Manganous ions in the water before the appearance of ferrous ions indicates that manganic precipitates in the oxidized mud surface are reduced at a higher potential than the ferric compounds.

Oxidation and reduction in soils seem to be regulated by the prevailing conditions of air circulation and by the oxygen consumption of microorganisms. Waterlogging of a soil causes a decided decrease in the redox potential (Stobe, 1956). The change from an oxidizing to a reducing state is associated with the appearance of reduced, and more soluble, iron and manganese and a lower pH. Presumably, ferric iron is reduced to the ferrous state and dissolved in the soil solution. In the presence of atmospheric or dissolved oxygen, iron is oxidized from the ferrous to the ferric state. Microbiological action on organic

matter produces carbon dioxide. The solution of carbon dioxide in water liberates hydrogen ions and brings into the soil solution increased quantities of exchangeable iron, manganese, aluminum, calcium, and magnesium. The formation of iron-bearing organic complex ions is an important factor in stabilizing iron in solution under oxidizing conditions at concentrations of iron that would otherwise precipitate as ferric hydroxide.

The normal redox potential of the system $\text{Fe}^{+++} + e \rightleftharpoons \text{Fe}^{++}$ is given as 0.68 v in $\frac{1}{8}$ M H_2SO_4 by Smith and Richter (1944); the value 0.77 is generally accepted as standard at pH 0 at 25°C (Bray and Hershey, 1934).

Since Puri and Sarup (1938) found the relation of pH to Eh in soils to be nearly linear, they believe that the measurement of redox potentials in soils does not give results of any greater accuracy than could be obtained by pH measurements.

Knaysi and Dutky (1936) found that in a bacterial-culture medium of ordinary meat infusion broth, the strictly aerobic bacterium *Bacillus megatherium* grows almost entirely on and near the surface where oxygen is most plentiful; however, it may be induced to grow throughout the medium or even at the bottom by an increase in the redox potential of the medium. Redox potential thus may affect bacterial growth distribution and location in lake bottoms.

Ignatieff (1941), Lipmann (1937), and Moore (1914) point out that light affects the reduction of ferric iron.

Redox-potential effects are discussed by the following: Allison and Scarseth, 1942; Baas-Becking and Wood, 1955; Baas-Becking and others, 1956; Baas-Becking and others, 1957; Belcher and others, 1955; Blagovidov and others, 1957; Bloomfield, 1952; Broadbent, 1954; Copeland, 1957; Eristavi, 1953, 1955; Gantimurov, 1952; Gorbunov, 1957; Hewitt, 1957; Hutchinson, 1957; Kramli and others, 1954; Melnichenko, 1954; Prokhorova, 1957; Rabotnova and others, 1955; Remezov, 1929; Schoeller, 1955; Shkolnik and Steklova, 1954; Trojanowski, 1954; Varkov and others, 1950; Warburg and others, 1933; Weart and Margrave, 1957; Yamasaki, 1952.

The effect of other factors in the reduction of ferric iron, such as waterlogging of soil and formation of complex iron compounds, is discussed by: Bezier, 1943; Brown, 1934; Deevey, 1941; Hill and Michaelis, 1933; Hirsch and Ruter, 1926; Hutchinson and others, 1939; Izgarishev and Turkovskii, 1929; Limanowski, 1932; Mason, 1952; Nitschmann, 1938; O'Meara, 1937; Osugi and others, 1934; Pear-sall, 1950; Puri and Sarup, 1938; Remezov, 1929, 1930; Robinson, 1930, 1951; Serdobolskii, 1950a, 1950b; Serdobolskii and Shavrygin, 1950; Shcherbina, 1939; Willis, 1936.

ORGANISMS UTILIZING IRON IN METABOLISM

The following organisms apparently require iron for normal metabolism according to the literature:

FUNGI

[See also iron bacteria]

Iron-using fungi are as follows:

- Aspergillus niger* (Bertrand and De Wolf, 1956; Bortels, 1927; Gollmick, 1936; Javillier and Sauton, 1911; Linossier, 1910; Lutman, 1929; Steinberg, 1919, 1920, 1935a, 1936)
oryzae (Nehira, 1955; Yamagata and others, 1955)
Cercospora nicotinae (Steinberg, 1950)
Fusarium lini (Saraswathi-Devi, 1955)
moniliforme (Saraswathi-Devi, 1955)
orthoceras (Saraswathi-Devi, 1955)
oxyспорum (Saraswathi-Devi, 1955; Steinberg, 1950)
poae (Saraswathi-Devi, 1955)
scirpi (Saraswathi-Devi, 1955)
udum (Saraswathi-Devi, 1955)
vasinfectum (Mundkur, 1928; Saraswathi-Devi, 1955)
Fusarium sp. (Saraswathi-Devi, 1955)
Helminthosporium sativum (Peterson and Katznelson, 1954)
Neurospora crassa (Nicholas and Commissiong, 1957)
Pythium irregulare (Steinberg, 1950)
Rhizoctonia solani (Steinberg, 1950)
Sclerotium rolfsii (Steinberg, 1950)
Streptomyces griseus (Kramli and others, 1954)
Thielaviopsis basicola (Steinberg, 1950)
Trichophyton mentagrophytes (English and Barnard, 1955)
rubrum (English and Barnard, 1955)
Ustilago sphaerogena (Garibaldi and Neilands, 1955)
Molds (Harley, 1940; Humfield and Sugihara, 1952; Metz, 1930; Muth and Voigt, 1929; Sauton, 1911)

Some soil fungi grow at pH environments approaching 1.0 (Truog, 1946).

BACTERIA

[See also iron bacteria]

Iron-using bacteria are as follows:

- Aerobacter aerogenes* (Waring and Werkman, 1943)
indologenes (Waring and Werkman, 1942, 1943)
Azotobacter chroococcum (Greaves, 1933; Krzemieniewski and Kovats, 1937)
vinelandii (Burk and others, 1931; Horner and Burk, 1934)
sp. (Burk and others, 1932)
Bacillus prodigiosus (Bortels, 1927)
bruntzii (Marchal and Adams, 1956)
Brucella abortus (Evenson and Gerhardt, 1955)
Candida guilliermondia (Enari, 1955)

- Chlorobium limicola* (Larsen, 1953)
thiosulfatophilum (Larsen, 1953)
Clostridium tetani (Feeney and others, 1943; O'Meara, 1937)
Escherichia coli (Waring and Werkman, 1943)
Ferrobacillus ferrooxidans (Leathen and others, 1956)
Flavobacterium halobium (Brown and Gibbons, 1955)
Klebsiella pneumoniae (Waring and Werkman, 1942, 1943)
Mycobacterium phlei (Tysarowski, 1955)
tuberculosis (Cohn and others, 1954; Henley, 1925; Minami and others, 1955; Polster and others, 1953)
Pseudomonas aeruginosa (Waring and Werkman, 1942, 1943)
cattleyae (Marchal and Adams, 1956)
cutirubra (Brown and Gibbons, 1955)
salinaria (Brown and Gibbons, 1955)
Rhizobium meliloti (Thorne and Walker, 1936)
trifolii (Thorne and Walker, 1936)
Rhizobium spp. (Virtanen and others, 1947)
Rhodopseudomonas sp. (Lascelles and Cooper, 1955)
Sarcina littoralis (Brown and Gibbons, 1955)
Serratia marcescens (Novelli, 1953; Waring and Werkman, 1943; Williams and others, 1956)
Thiobacillus denitrificans (Baalsrud and Baalsrud, 1954)
ferrooxidans (Bryner and others, 1954; Temple and Colmer, 1951)
thiooxidans (Bryner and others, 1954)
Bacilliform bacterium (Brussoff, 1918)
Bacteria (Anderson, 1940; Kaserer, 1910; Lasseur and others, 1944; Waring and Werkman, 1942)
Bacteria producing blackbeet disease (Cameron, 1934)
Hemophilic bacteria (Lwoff, 1933b)
Intestinal flora (Stern and others, 1955)
Marine bacteria (MacLeod and Onofrey, 1956)
Marine sulfate-reducing bacteria (Kimata and others, 1955)
Soap-discoloring bacteria (Hollo and Gorog, 1954)

PROTOZOA

Iron-using protozoa are as follows:

- Chilomonas paramecium* (Bowen, 1940; Hutchens, 1940)
Paramecium candatum (Shoup and Boykin, 1931)
Pleurobrachia sp. (Cooper, 1939)
Trypanosoma sp. (Lwoff, 1933a)

PHAGES

The following workers have reported iron-using phages:

- Bortels, 1953; Spizizen, 1943

VIRUSES

Iron utilization by virus is also known.

- McIntire and others, 1941

YEASTS

Although yeasts are fungi they are listed separately in this report.

- Bortels, 1927; Elvehjem, 1931; Kiene, 1940; Malkov, 1934

ALGAE

Iron-using algae are as follows :

Blue-green

- Anabaena lemmermanni* (Guseva, 1937b)
- Aphanizomenon flos-aquae* (Guseva, 1937b)
- Coclosporium* sp. (Guseva, 1939)
- Lyngbia confervoides* (Kinne-Diettrich, 1955)
- maincula* (Kinne-Diettrich, 1955)
- Microcystis aeruginosa* (Gerloff and Skoog, 1957)
- Oscillatoria agarhdia* (Skadovskii and others, 1955)
- rubescens* (Edmondson and others, 1956)
- Phormidium ambiguum* (Umemoto and Mifune, 1953)
- Blue-green algae in general (Slobodchikov and Stroikina, 1953; Vouk, 1936) ; (Gerloff, written communication, 1955)

Brown (ocean kelp)

- Macrocystis integrifolia* (Wort, 1955)
- Nereocystis luetkeana* (Wort, 1955)

Desmids

- Desmidiaceae (Hofler, 1925)

Diatoms

- Asterionella formosa* (Cooper, 1935; Guseva, 1937b)
- japonica* (Goldberg, 1952)
- Chaetoceras curvisetius* (Koblents-Mislake, 1955)
- Nitzschia closterium* (Harvey, 1933; Koblents-Mislake, 1955)
- Diatoms in general (Correns, 1941; Gayral, 1954; Harvey, 1937, 1939; Thompson and others, 1932)

Green

- Chlamydomonas* sp. (Lackey, 1939)
- Chlorella pyrenoidosa* (Knauss and Porter, 1954; Myers and others, 1951; Walker, 1954)
- vulgaris* (Mayer, 1952; Mayer and Gorham, 1951)
- sp. (Hopkins, 1930, 1935; Hopkins and Wann, 1926, 1927; Reisner and Thompson, 1956; Schwartz, 1956)
- Cladophora glomerata* (Blum, 1953, 1956)
- Euglena* sp. (Lackey, 1939)
- Ulothrix* sp. (Lackey, 1939)
- Green algae in general (Roberg, 1932; Slobodchikov and Stroikina, 1953)

General

- (Blum, 1956; Cooper, 1935; Ellis, 1925; Fogg, 1952; Gayral, 1954; Gorham and Gorham, 1955; Goryunova and Nasonova, 1955; Guseva, 1937a; Hutchinson, 1941; Koblents-Mislake, 1955; Lackey, 1956; Levring, 1957; Messikommer, 1956; Moore, 1950; Murray, 1908; Novozhilova, 1955; Ono, 1900; Pond, 1903; Prescott, 1956; Pringsheim, 1934; Riley, 1940; Roberg, 1932; Schroder, 1939; Slobodchikov and Stroikina, 1953; Smith, 1950; Uspenski, 1927; Villeret, 1954a, 1954b, 1954c, 1954d, 1954e; Voronkov 1953; Yoshimura, 1936)

MICRO-ORGANISMS, GENERAL

The following workers have reported use of iron by general micro-organisms:

Neilands, 1957; Osugi and Aoki, 1938b; Scouller, 1932

FLOWERING PLANTS

Iron-using flowering plants are as follows:

Aquatic, rooted (also see crop plants, rice)

Oryza sativa (Juliano and Aldama, 1937; Richter, 1926; Tullis and Cralley, 1936)

Potamogeton foliosus (Harper and Daniel, 1934-35)
spp. (Nierenstein, 1945)

Typha latifolia (Harper and Daniel, 1934-35; Lackey, 1939)

Watercress (Escudero and others, 1944; Gorham and Gorham, 1955)

General (Ward and Whipple, 1918)

Aquatic, unattached

Lemna major (Fly, 1935; Yoshimura, 1950)

minor (Steinberg, 1941, 1943, 1946)

polyrhiza (Olsen, 1930; Saeger, 1937)

sp. (Clark, N. A. 1936; Hopkins, 1935)

Aquatic, general

(Bowman, 1956; Butcher, 1933; Curry and Wilson, 1955; Dupont, 1954; Einsele, 1936a, 1936b; Lewis and Goldberg, 1954; Margalef, 1954; Mayer and Gorham, 1951; Moore, 1952; Olsen, 1950; Olson and others, 1941; Pond, 1903; Prescott, 1956; Roelofs, 1940; Rozenberg and Mefedova, 1956; Skadovskii and others, 1955; Villeret, 1954e)

Crop

Truog (1946) has shown that a soil pH of 6.5 is very favorable for absorption of iron by most crop plants.

Alfalfa (Dalle, 1946)

Apple (Milad, 1939)

Asparagus (Sheets and others, 1941)

Cabbage (MacGillivray and others, 1942; Sheets and others, 1941)

Carrots (MacGillivray and others, 1942)

Clover (Brewer and Carr, 1927)

Corn (Allyn, 1927; Arndt, 1922; Brewer and Carr, 1927; Brown and Holmes, 1955; Eckstein and Jacob, 1929; Hoffer and Carr, 1920; Jacobson, 1945; Maze and others, 1912; McGeorge, 1939; Olsen, 1938; Sayre, 1930; Scharrer and Schropp, 1933)

Lettuce (Harward and others, 1955; Sheets and others, 1941; Twyman, 1951)

Mustard greens (Abbott and Ahmann, 1940)

Oats (Mayer, 1892; Twyman, 1951)

Parsley (Randoin and Fournier, 1947)

Pea (Brown and Possingham, 1957; Possingham and Brown, 1957)

Pear (Bennett, 1928; Jacobson, 1945; Milad, 1939)

Pineapple (Gile and Carrero, 1916a; Sideris and Young, 1956)

Potatoes (MacGillivray and others, 1942)

Quince (Rodriguez and Abadia, 1956)

Rice¹ (Aiyar, 1946; Gile and Carrero, 1914, 1916a, 1916b; Kapp, 1934, 1938; Milad, 1939; Richter, 1926; Tullis and Cralley, 1936; Ueda, H., 1956; Ueda, K., and others, 1956; Willis and Carrero, 1921)

Soybean (Brown and Holmes, 1955; Marsh and Shive, 1925)

Spinach (MacGillivray and others, 1942)

Sugar beet (Whatley and others, 1951)

¹ Rice grows as an emergent aquatic plant. (Also see flowering plants, aquatic, rooted).

Sugarcane (Martin, 1935)

Tobacco (Jacobson, 1945; McMurtrey, 1933, 1952)

Tomato (Porter and Thorne, 1955; Twyman, 1951)

Tung (Dickey, 1942)

Turnip greens (Abbott and Ahmann, 1940)

Wheat (Barnette, 1921; Barnette and Shive, 1923; Brewer and Carr, 1927; Brown and Holmes, 1955; Greer and others, 1952; Singh, 1955, 1956)

Winter squash (MacGillivray and others, 1942)

General (Guelbenzu, 1951; Harris, 1952; Hewitt, 1950, 1953; Martin, 1935; McGeorge, 1951; Nicholas, 1949; Peterson and Elvehjem, 1928; Porter and Thorne, 1955; Randoin and Le Gallic, 1942; Remington and Shiver, 1930; Wallace, 1954; Wiebosch, 1947; Winsor, 1939)

Other than crop

Artemisia sp. (Susplugas and others, 1951)

Azalea spp. (Carnes, 1954)

Gardenia sp. (Baudenistel, 1957)

Hydrangea sp. (Atkins, 1923)

Rhododendron spp. (Carnes, 1954)

Xanthium pensylvanicum (Smith and others, 1957)

Roses (Carnes, 1954)

Sunflower (Carnes, 1954; Jacobson and Oertli, 1956)

Trees (Arnaud, 1919)

ANIMALS

The following workers have reported use of iron by animals:

Grazing on iron-deficient vegetation

Aston, 1934; Wind, 1951.

Nongrazing (other than man)

Davis and Loosli, 1954; Free, 1940; Galtsoff, 1934; Inoue, 1932; Kosugi and Umeda, 1931; Newell and McCollum, 1931; Parks and Rose, 1933; Supniewski, 1951; Unti, 1943.

Man

Adams and others, 1950; Chittenden, 1896; Elvehjem and others, 1927; Hendrych and Klimesch, 1935; Johnston and others, 1950; Lanyar and others, 1933; Lederer and Bogaert, 1939; Lindquist, 1949; Lintzel, 1929; McCance and Widdowson, 1943; Palit and others, 1944; Perosa and others, 1951; Piery and others, 1938; Sharpe, 1948; Steinkamp and others, 1955; Will and Vilter, 1954.

ORGANISMS, GENERAL

The following workers have reported use of iron by general organisms:

Antognini, 1954; Bear, 1954; Blunno, 1911; Boiret and Paturel, 1892; Bollmann and Schwanitz, 1957; Bonner, 1950; Frisbie, 1953; Granick, 1954; Ignatieff, 1941; Ingalls and Shive, 1931; Jenny and Overstreet, 1939; Keilin and Mann, 1944; Kliman, 1937; Loew, 1938; Masoni, 1914; Monnier and Kuczarski, 1917; New Jersey Agricultural Experiment Station, 1928; Nikitin, 1950; Olson, 1947; Parsche, 1940; Pirrie, 1950; Rankama and Sahama, 1952; Robinson, 1951; Stoklasa, 1898; Wallbach, 1932; Wann, 1930.

IRON BACTERIA

[See also bacteria and fungi]

The iron bacteria constitutes a heterogenic group of organisms that are difficult to define. For example, Naumann (1928a) defined the group as all organisms that can precipitate or dissolve iron. However, the literature has described numerous organisms which dissolve and use iron in metabolism. Cholodnii (1928) has a definition more restrictive but apparently inadequate and even erroneous. Cholodnii believed the term "iron organism" should be restricted to organisms that respire ferrous oxide. The literature shows that on the spot, organic material rather than ferrous oxide is essential for the respiration of these organisms. The term "iron bacteria", as generally considered, includes many organisms that are not bacteria.

The following list of characteristics shows that the iron bacteria belong to a transition group, the actinomyces, which are fission fungi or thread fungi having both bacterial and fungal characteristics:

*Bacterial characteristics*Thallus width 0.5–1.2 μ

Some bacteria form only a rudimentary mycelium.

Relative primitive state of the nucleus

Fungal characteristics


Reproduction by aerial budding out of successive conidia by abstriction as for example, yeast of ascomycetes.

Ability to form a true coenocytic mycelium.

Normal ability or tendency to form a mycelium under conditions suitable for active vegetative growth.

Lackey and Wattie (1940) were able to isolate only the iron bacterium *Sphaerotilus natans* from Ohio River sewage water although these field collections did show morphological differences. Pringsheim (1949) believed the many morphological field manifestations of the same iron bacterium to be responsible for the many synonyms in the literature. Thus, he found *Cladothrix dichotoma* Cohn, *Leptothrix ochracea* Kutzing, *Chlamydothrix ochracea* (Kutzing) Migula, *Chlamydothrix sideropons* Molisch, and *Clonothrix fusca* Roze to be synonyms of *Sphaerotilus natans* Kutzing. (See fig. 13 for diagrams of different iron bacteria.)

In spite of considerable overlap, classification according to morphological form is useful for a fast field identification. Accordingly, iron bacteria are classified by genera, using the original worker's nomenclature. However, in classifying these organisms as iron bacteria, it must be remembered that many precipitate or dissolve iron in addition to other elements, such as manganese and sulfur. Generally, ferric hydroxide is deposited only in the thick sheathlike envelope of living organisms (California State Water Pollution Control Board, 1952; Reilhes, 1944), but it may be deposited elsewhere. Iron bacteria



Fluffy bits of floating or immersed cottonlike material often may be found in sewage-contaminated streams in early summer or late autumn

SPHAEROTILUS




Parallel threads with a common mucilaginous covering




Most widely distributed of all iron bacteria. Predominate in open, shallow water exposed to direct sunlight. Stiff rods usually lack cell contents

LEPTOTHRIX




Semimature
CLADOTRICH



Normally attached at one end to some foreign object such as a particle of dirt or a stone granule

CRENOTHRICH



Young



Old

A spirally twisted thread possessing no sheath or mucilaginous envelope; predominate in darkened waters in conduits and tunnels

GALLIONELLA



SPIROPHYLLUM

FIGURE 13.—Diagrams of different iron bacteria (after Ellis, 1919).

frequently act upon solutions of relatively low (less than 0.5 ppm) iron concentration. Deposits of ocher and bog ore made up of minute tubes of iron oxide are produced by them; the mucilaginous envelopes of the bacteria prevent the escape of the ferric hydroxide. In consequence, the iron begins to accumulate round the tubercle. The bacteria thus have an extraordinary capacity for storing iron and manganese on the sheath; the thickness of the encrusted sheath is sometimes 12 or more times that of the original filament (Ellis, 1919).

One of the earliest and most striking examples of the effect of iron bacteria is that recorded in the Book of Exodus (VII: 20-25) circa 500 B.C. In the account of 1 of the 7 plagues of Egypt, the water in the Nile River was "turned to blood," the fish died, the river became foul, and the Egyptians could not drink the water. These conditions probably were induced by the sudden growth of microscopic organisms. About 2,400 years later, in 1887, history repeated itself at Rotterdam, Holland, and the water became undrinkable (Yust, 1951); the walls of the reservoirs became thickly encrusted with mussels, which, in turn, became densely coated with $\frac{1}{4}$ - to $\frac{1}{2}$ -inch covering of filamentous iron bacteria that had entered the service mains. Ten years earlier, in 1877, the service pipes of Berlin, Germany, were choked completely by iron-bacteria deposits (Yust, 1951), and all the water mains had to be renewed.

Because *Sphaerotilus natans* and other typical iron bacteria develop only in an approximately neutral environment (Stokes, 1954) in which chemical oxidation of ferrous iron is rapid and extensive, the function of the bacteria in oxidation becomes difficult to determine. (California State Water Pollution Control Board, 1952; Glickhorn, 1920). Ferrous ions are readily oxidized in natural surface waters to insoluble ferric hydroxide. These precipitates tend to agglomerate, flocculate, and settle or be adsorbed on surfaces; hence, the concentration of iron in well-aerated waters is seldom high. Natural iron removal is fortunate as a concentration of more than 0.30 ppm is objectionable in waters for public supply. Probably many of the lower bacteria as well as thread bacteria function permanently or temporarily as iron bacteria (Ellis, 1919).

As cited in the literature the following organisms may be considered as iron bacteria according to popular usage of the phrase:

Manganese-metabolizing:

Anthophysa vegetans (Pringsheim, 1946)

Bacterium precipitatum (Kalinenko, 1946)

Bikosoeca petiolata (Pringsheim, 1946)

Crenothrix fusca (Thiele, 1928; Wasser, 1938)

sp. (Dienert, 1938)

Gallionella ferruginea (Thiele, 1928; Wasser, 1938)

Leptothrix echinata (Beger, 1935)

ochracea (Lieske, 1919; Thiele, 1928; Wasser, 1938)

sp. (Dienert, 1938)

Pseudomonas ferrugineum (Pillai, 1938)

Siderocapsa sp. (Dienert, 1938; Wasser, 1938)

Siderodendron manganiferum (Pringsheim, 1946)

Siphomonas fritschii (Pringsheim, 1946)

Gel-forming bacteria : (Beger, 1935; Beger and Haase, 1941)

Manganese bacteria : (Baer, 1933; Hallsworth and Costin, 1953)

Sulfur-metabolizing:

Because sulfate-reducing bacteria seem prominent in causing the aqueous solution of otherwise insoluble iron, the environmental conditions favorable to the growth of this group of bacteria (Ivanov, 1956) are listed: (a) neutral pH (about 7); (b) absence of oxygen; (c) a source of organic nutrients; (d) a source of sulfates.

Chlorobium limicola (Larsen, 1953)

thiosulfatophilum (Larsen, 1953)

Spirophyllum sp. (Meehan and Baas-Becking, 1927)

Thiobacillus ferrooxidans (Temple and Colmer, 1951)

Toxothrix sp. (Meehan and Baas-Becking, 1927)

Vibrio desulphuricans (Butlin and others, 1952)

Anaerobe (Austen, 1931)

Marine bacteria (Francis-Boeuf, 1948)

Sulfur bacteria (Kimata and others, 1955; Rudolfs, 1922; Starkey, 1945;

Starkey and Wight, 1943)

Organic iron metabolizing:

Cladothrix dichotoma (Ellis, 1921)

Crenothrix polyspora (Ellis, 1921)

sp. (Whipple and others, 1927)

Gallionella ferruginea (Ellis, 1921)

Leptothrix ochracea (Ellis, 1921; Kalinenko, 1940)

Pseudomonas ferrugineum (Pillai, 1938)

Spirophyllum ferrugineum (Ellis, 1921)

Microorganisms, general (Allison and Scarseth, 1942; Beger and Haase, 1941; Betremieux, 1951, 1954; Starkey, 1945; Thornton, 1956; Yoshimura, 1936)

Calcium-metabolizing:

Achromatium oxaliferum (Lackey, written communication 1959)

Bacterium precipitatum (Kalinenko, 1946)

Pseudomonas ferrugineum (Pillai, 1938)

Other-metabolizing:

Crenothrix fusca (Wasser, 1938)

polyspora (California State Water Pollution Control Board, 1952;

Duchon and Miller, 1948)

Gallionella ferruginea (California State Water Pollution Control Board, 1952; Beger, 1937; Volkova, 1939; Wasser, 1938)

Leptothrix ochracea (California State Water Pollution Control Board, 1952; Beger, 1937; Glickhorn, 1920; Volkova, 1939; Wasser, 1938)

trichogenes (Beger, 1937; Gleen, 1950)

Siderocapsa sp. (Wasser, 1938)

Sphaerotilus natans (Stokes, 1954; Wasser and others, 1937)

Trachelomonas sp. (Glickhorn, 1920)

Iron bacteria, general (Banfi, 1952; Baylis, 1925; Betremieux, 1954; Dementiev, 1941; Fujihara, 1925; Haase, 1940; Harmsen, 1938; Hutchinson, 1941; Kliman, 1937; Kurzweil and Exner, 1954; Naumann, 1928a; Pringsheim, 1946; Scott and Brandly, 1933; Sorokina, 1938; Tillmans and Klarmann, 1924; Varkov, and others, 1950; Waring and Werkman, 1943; Winogradsky, 1922)

Control:

Methods of controlling iron bacteria have been suggested by the following investigators:

Anthrophysa vegetans (Cataldi, 1937)

Asellus aquaticus (Holland, 1956)

Crenothrix polyspora (Duchon and Miller, 1948)

sp. (Grime, 1945; Hale, 1949)

Leptothrix ochracea (Cataldi, 1937)

Iron bacteria, general (Baer, 1933; Beger and Haase, 1941; Derby, 1956; Ellis, 1921; Gleen, 1950; Griffin, 1937; Haase, 1940; Harvey, 1939; Osugi and others, 1935; Pronin, 1933; Thiele, 1928; Tillmans and Klarmann, 1924; Volkova, 1939)

Fungus slimes (Hale, 1949)

Table 1 summarizes the many organisms that require iron for normal metabolism. Large forms of animal life are excluded from the table.

TABLE 1.—Organisms that require iron for normal metabolism

Organism	Number of genera	Number of species
Algae:		
Blue-green.....	6	7
Desmids.....	(¹)	—
Diatoms.....	3	4
Green.....	5	3
Other.....	3	4
Bacteria.....	20	25
Flowering plants (aquatic):		
Soil-rooted.....	3	3
Unattached.....	1	3
Flowering plants (land):		
Crop.....	25	—
Noncrop.....	7	—
Fungi.....	12	23
Iron bacteria:		
Calcium-metabolizing.....	2	2
Manganese-metabolizing.....	10	10
Organic iron metabolizing.....	6	6
Sulfur-metabolizing.....	5	4
Other-metabolizing.....	6	5
Phages.....	(¹)	—
Protozoa.....	4	4
Viruses.....	(¹)	—
Yeasts.....	(¹)	—

¹ 1 or more.

Thus, studies made by many investigators of the normal metabolism of numerous organisms affirm the essential function of iron in vital life processes.

MECHANICS OF IRON CONCENTRATION AND DEPLETION

LICHEN EFFECTS

Not only do lichens take certain chemical elements from the rocks, but after their death and decay, they release organic acids that increase the chelating or solvent power of the natural water (Schatz and others, 1956). The solubility of iron is much greater in the presence of these organic acids.

The brown or black shiny crust on rocks known as desert varnish consists mainly of oxides of iron and manganese. Lichens probably are important in contributing to such surface-crust deposition (Emmons and others, 1955). Also, lichens may be concentrators of iron (Yarilova, 1947).

HYDROPHYTE (OTHER THAN BACTERIA AND FUNGI) EFFECTS

The chlorophyll-bearing water plants are important in supplying iron to and removing iron from bodies of water. Algae, characteristically haploid (gametophyte) organisms generally lacking a mineral-conducting system, extract iron and other mineral salts directly from the water medium (Gessner, 1956; Glenck, 1954; Gloyna and others, 1955; Henderson and others, 1954; Smith, written communication, 1956; Steel and Gloyna, 1955; Walker, 1956; Yoshii and Watabe, 1956). Smith, 1950, has an excellent discussion of haploid-diploid variability in the algae. Soil-rooted aquatic plants, characteristically diploid (sporophyte) organisms generally having an excellent mineral-conducting system, extract iron and other mineral salts directly from the water-immersed soil medium; thus, they live aerobically in anaerobic growth situations (Crafts, 1954; Lowenhaupt, 1955; Marei, 1955; Pond, 1903; Prescott, 1956). Normally free-floating aquatic plants, whether haploid or diploid, absorb minerals directly from their water environment. A diagram of the relative importance of water-mineral and soil-mineral extraction during the life cycles of the four general types of plants is given in figure 14. Five features that characterize aquatic plants and emphasize their large use of iron are as follows:

1. Large proportional amount of chlorophyll-bearing tissue which results from (a) large expansion of plant body surface by lengthening and subdivision of plant parts, (b) assimilatory region, generally confined to the increased epidermal area. The work of Leibich (1941) cited by Bonner (1950) shows that as




FIGURE 14.—Relative importance of the gametophyte (single line of the life-cycle circle representing water mineral absorbing) and sporophyte (stippled part of the life-cycle circle representing soil mineral absorbing) generations in the four major groups of aquatic plants (adapted from Milne and Milne, 1952).

much as 82 percent of the iron of spinach leaves occurs in the chloroplasts. Of this, at least four-fifths was found in organic combination. It seems logical to assume the same would be true for aquatic-plant chloroplasts.

2. Reduced organic iron compounds are easily extracted from soils devoid of oxygen. Large air-conducting spaces, through which oxygen for root respiration is sent from the leaves, in frequently cockscrew-shaped water roots are present. These air-conducting spaces enable the roots to penetrate far into water-saturated un-aerated mud (Fujii and Tanaka, 1955). These anaerobic situations have a low redox potential. Land-plant soil roots would suffocate (Gausman and Struchtemeyer, 1957; Jackson, 1956) under similar undrained growth conditions because in land plants, oxygen for root respiration is supplied to the root directly from the air in the soil. The results is that aquatic plants readily and effectively, through growth and disintegration of their dead bodies, transfer or move iron from the soil to the water medium. Figure 15 illustrates the root and leaf structures of aquatic and land plants.
3. The xylem which is usually considered to be tissue which conducts inorganic minerals frequently may be deficient as in Sago pondweed (Eames and MacDaniels, 1925). Aquatic plants have a well-developed phloem, or organic food-conducting tissue, through which reduced organic iron compounds may readily travel (Arber, 1920; Arisz and Schrender, 1956; Eames and MacDaniels, 1925; Kramer, 1957; Schimper, 1903).
4. The plants are characterized by (a) a high ash content, (b) a high protein content (Dubois, 1955), and (c) a very low crude fiber content. The legume hays are the only commonly used forages which can approximate aquatic plants in these constituents (Gortner, 1934).




FIGURE 15.—Morphological (and therefore biochemical functional) comparison between leaf midrib and root structure of rice and comparable parts of typical aquatic and land plants. Rice plants have a typical aquatic plant structure as indicated by the large "air cells."


5. Percentage of iron in ash is high. For the common water plant *Elodea*, iron has been reported as 2.56 percent of the ash (Bonner, 1950).

Aquatic plants further reduce the quantity of suspended and (or) colloidal iron by (a) preventing mechanical silting and erosive wave action and (b) chemical clarifying action on water containing suspended matter (De Gruchy, 1938).

As excellent oxygenators of water during daylight hours, aquatic plants tend to remove the ferrous iron in aqueous solution.



Thus, both aeration and photosynthesis in water-growth situations cause excess iron to flocculate and settle to the mud bottom. The principle of most iron-removal processes involves the neutralization of hydrogen ions and the increase in oxygen (Calise and Dietz, 1955; Klyachko, 1956; Weston, 1936). Conversely, during the night when photosynthesis is low and respiration is rapid, aquatic plants will return some iron to solution (Marcellin, 1956; Newell, 1957).



However, in open-water bodies such as streams and lakes, the hydrogen-ion concentration probably cannot be raised enough by this effect to greatly increase the iron concentration.

The removal of dissolved organic and inorganic constituents in water by freezing has been investigated by Mortimer (1942). He found that 94 percent of dissolved salt, all humus coloring matter, and a large part of dissolved gas were removed from surface water by freezing. Water freezing thus resulted in an increase of the mineral content in the water immediately under the ice.

Tannins, which may form stable complexes with iron and other metallic ions are found in several parts of the aquatic plant, especially in epidermal cells of leaves, in the cortical tissues of stems, and in the walls or vacuoles of cells.

The plant *Enteromorpha*, which can exist in fresh water as well as in water more mineralized than sea water, has been found by Baas-Becking and Mackay (1956) on intertidal flats. It produces highly reducing acidic substances in the dark, such as dimethyl sulfonium compounds. This reducing environment probably causes some iron to go into solution. In the light, some of these compounds are either reabsorbed or destroyed; thus precipitation of iron results.

The simpler aquatic plants, the algae, may be attached or unattached to the substratum. (See p. 114 for discussion of their iron assimilation.) Some algae, such as the stoneworts among the fresh-water algae and some seaweeds in marine algae, apparently have effective conduction systems and, absorb iron from the mud substratum. However, this type is the exception rather than the rule.

The more complex flowering plants may be rooted or unattached to the substratum. Those unattached absorb iron directly from the water bathing the organism or from the water upon which the organ-

ism floats. (See p. 114 for discussion of the iron assimilation of attached flowering plants.)

Because aquatic plants generally occur in bodies of surface water, and frequently in great abundance their relationships to iron in water are emphasized. Because of proportionately high water and low cellulose contents, solid materials in the water-plant structure disintegrate rapidly upon death. Thus, when living, aquatic plants assimilate matter that may be used as food; and in death, they yield important salts and organic substances to the water. Tannin and other organic compounds that may be yielded by disintegration of dead plant material are important in holding the iron in solution.

Figure 16 shows some of the processes involved in the circulation of iron between soils and water, and the interrelationships between iron in dead organisms and iron used by the living. Both aerobic and anaerobic bacteria release and thereby make available iron to the new crop of plant growth. Animals either directly or indirectly feed on the plant organisms and by assimilation transfer iron to their bodies.

Iron must be present in plants for production of chlorophyll. Upon death and disintegration of the plant material, iron is released into the water. During the following growing season, which may be immediate, iron would again be taken from the substratum by the growing plant. Thus, there is an annual transfer of iron from the substratum to the water body. Under aerobic conditions, iron is oxidized and precipitated. Apparently, anaerobic bacterial disintegration is important in releasing iron from its organic complexes in dead plant tissue. Practically, anaerobic conditions in water are characterized by reduction of nitrogen and sulfur compounds and the appearance of methane as well as absence of oxygen (Baas-Becking and Kaplan, 1956; Baas-Becking and Mackay, 1956; Kufferath and Schmitz, 1955). In oxygen-deficient environments, sulfate may be reduced to hydrogen sulfide, which has an offensive odor of putrefaction, precipitates usually as black metallic sulfides, and may be toxic to some forms of life. In a strongly reducing environment, protein and its primary cleavage products are broken down by hydrolysis rather than by oxidation. Such breakdown causes foul-smelling mercaptans and amines such as accompany the decomposition of flesh.

Iron is transported in the phloem tissue of higher plants and is concentrated in the mesophyll or in the cambium (Kliman, 1937). Regions of plant tissue in which iron and other metallic salts tend to concentrate are illustrated in figure 17. Most active salt accumulation in the higher aquatic plants is in the meristem zone, for example,





FIGURE 16.—Transverse portion of lake. Some phases of the iron cycle in A, by the macrobiota, and in B, by the microbiota, which affect the presence of iron in natural water.

root, shoot, and leaf base (Steward, 1948). Each lateral bud as a potential growing region is a potential region of salt accumulation.

Several hydrophytes are important concentrators of iron. Iron concentration of 26,000 times that of the immersing water medium has been recorded in some seaweeds (Oy, 1940). The following aquatic plants are reported to be concentrators of iron:

*Hydrophyte concentrator and reference**Cladophora* sp. (Whipple and others, 1927)*Chlorella* sp. (Mayer and Gorham, 1951)*Crenothrix* sp. (Whipple and others, 1927)*Fucus vesiculosus* (Oy, 1940)*Iridaea* spp. (Read and Gow, 1927)*Laminaria digitala* (Oy, 1940)*L. japonica* (Read and Gow, 1927)*L. religiosum**Microcystis aeruginosa* (Gerloff, 1955)*Nostoc flagelliforme* (Read and Gow, 1927)*Oedogonium* sp. (Ellis, 1919)*Potamogeton* spp. (Read and Gow, 1927)*Rhodymenia* spp. (Read and Gow, 1927)*Sargassum siliquastrum* (Read and Gow, 1927)

Algae in general (Smith, written communication, 1956)

Aquatic plants, general (Mayer and Gorham, 1951; Speirs, 1954)

Desmids (Ellis, 1919; Hofer, 1925)

Diatoms (Thompson and others, 1932)

Marine brown algae (Black and Mitchell, 1952)

Watercress (Escudero and others, 1944)

Zygnemaceae (Ellis, 1919)

General (Harder, 1919; Naumann, 1928b)




FIGURE 17.—Regions of dicotyledon (left) and monocotyledon (right) plants in which iron and other metallic salts tend to concentrate. These are regions of high metabolic activity.

A lush growth of *Cladophora* was recently found growing in a Colorado spring that yielded water high in iron content. Microscopic examination revealed that filaments of this alga had heavy pellicle deposits of ferric hydroxide similar to those commonly found on the iron bacteria *Leptothrix*. A similar phenomenon has been reported by Rackow (1957) for *Chlorella pyrenoidosa* and *Ulothrix zonata*. Gerloff (written communication, 1955) reported the iron content of algae ranges from 164 to 543 ppm.

Oxygen has been mentioned as one of the elements that lowers the solubility or alters the availability of iron in water. It also is true (Gerb, 1956; Speller, 1922, 1951; Walker and others, 1907, 1908) that the rate of corrosion of metallic iron in natural water is almost directly proportional to the oxygen content for concentrations up to about 6 cubic centimeters per liter (according to Stoltenberg, 1949, 2 cc per liter). According to Nietsch (1956), water containing a dissolved-oxygen content lower than 4 cubic centimeters per liter tends to dissolve iron. Oxygen, which increases galvanic corrosion, is an active depolarizer at all temperatures, and even in an alkaline solution. The processes of corrosion are not covered in this report.

CHEMICAL ELEMENTS INTERFERING WITH USE OF IRON BY PLANTS

Several chemical elements, when in water, interfere with the use of iron by plants. Some of these elements may decrease the solubility of iron or act as oxidizing agents in causing ferrous iron to be changed to insoluble and unavailable ferric iron. Others may react in different ways to prevent normal assimilation of iron. Calcium content, which affects lake and land productivity of plants, is mentioned by Aiyar, 1946, 1948; Allyn, 1927; Bennett, 1928; Boischot and Durroux, 1950; Brown and Gibbons, 1955; Crawford, 1939; De and Basu, 1949; De Kock, 1955c; Demolon and Bastisse, 1938; Doyne and Morison, 1926; Gayral, 1954; Gayral and Panouse, 1954; Gile and Carrero, 1914, 1916b; Hasler, 1951; Hopkins and Wann, 1927; Iyengar and Subrahmanyam, 1935; Kuhnholz-Lordat, 1928; Masoni, 1914; Maze and others, 1912; McGeorge, 1951; McGeorge and Breazeale, 1956; Milad, 1939; Parsche, 1940; Robinson, 1951; Samish, 1954; Schachtschabel, 1942; Sidorin, 1914; Simakov and Isakova, 1933; Skeen, 1930; Steele, 1949; Stumm, 1956; Upadhyaya, 1955; Van der Spek, 1948; Vidal, 1938; Villeret, 1954e; Wallace and Ashcroft, 1956; Wallace and others, 1955; Wann, 1930; Wiebosch, 1947.

Copper in water acting as an oxidizing agent and causing ferrous hydroxide to form insoluble complexes (Puustjarvi, 1952) prevents

algae from using iron. These and other effects are mentioned by: Aiyar, 1946; Beaumont and Holland, 1935; Brown and Holmes, 1955; Cervenka, 1955; Coupin, 1898; Crabill, 1956; Curini-Galletti, 1937; De and Basu, 1949; Densch and Hunnius, 1924; Echave, 1937; Elvehjem, 1935; Erkama, 1949; Fattinger, 1950; Forster, 1953; Guseva, 1940; Hale, 1949; Harrison and Subrahmanyaayyar, 1917; Harward and others, 1955; Hood, 1948; Isizuka, 1940; Keilin and Mann, 1944; Koblents-Mislake, 1955; Korinek, 1936; Lahey and others, 1952; Lazarev, 1939; Magie, 1951; Malyuga, 1945; Maquenne and Demoussy, 1920; Marsais and Segal, 1938; Moore and others, 1957; O'Meara, 1937; Ono, 1900; Ornstein, 1935; Pika, 1956; Posnjak, 1926; Puustjarvi and Juusela, 1952; Quartaroli, 1932; Riley, 1939; Rippel, 1941; Saeger, 1937; Schmidt-Nielsen, 1938; Schwaibold and Fischler, 1931; Shkol'nik and Makarova, 1950; Voznesenskii and Nagatkin, 1934; Whatley and others, 1951; Williams, 1953; Willis, 1936; Willis and Piland, 1936, 1937.

Watson and Bollen (1952) found a definite inhibition in bacterial growth in the presence of a large amount of organic matter when the total copper concentration ranged from 11 to 30 ppm. However, no deleterious effect on bacterial growth was observed in controls containing 4 to 7 ppm in concentration of total copper and low organic matter content. Conditions unfavorable for growth would no doubt decrease the amount of iron brought into solution by bacteria.

Magnesium is reported by Aiyar (1948), De and Basu (1949), and Schachtschabel (1942) to interfere in iron metabolism of plants.

Manganese is reported by the following writers to inhibit iron utilization by plants: Aiyar, 1946; Beger, 1935; De and Basu, 1949; Guseva, 1937a; Johnson, 1916; Kapp, 1932a, 1932b; Kimata and others, 1955; Kitano, 1955; Kuhnoltz-Lordat, 1928; Magie, 1951; Martin, 1935; Ouellette, 1951; Rippel, 1923; Somers and Shive, 1942; Willis and Piland, 1937; Yoshimura, 1931c; Zapffe, 1931, 1933. Excessive manganese can produce chlorotic symptoms which indicate iron deficiency of plants.

Water high in oxygen usually does not permit more than minor traces of iron in solution. Aspects of this are discussed by: Beger, 1937; Bochkarev and Yanko, 1953a, 1953b; Brozek, 1949; California State Water Pollution Control Board, 1952; Chernovskaya, 1954; Correns, 1941; Dementiev, 1941; Dienert, 1938; Einsele, 1940a, 1940b; Fabrin, 1948; Gayral and Panouse, 1954; Griffith, 1955; Haase, 1940; Hutchinson, 1941; Hutchinson and others, 1939; Ingleson and Thomas, 1946; Ivlev, 1938; Juday and others, 1938; Kriegsmann, 1938; Malyuga, 1945; Mancha, 1936; Martin, 1935; Muller, 1950; Nietsch, 1956; Pearsall, 1950; Pillai, 1938; Rankama and Sahama, 1952; Saunders,

1938; Seeberg, 1937; Stoltenberg, 1949; Strakhov, 1948; Sugawara, 1934; Tanaka, 1953, 1954; Thompson and others, 1932; Thompson and Bremner, 1935b; Todt, 1950; Urbain and Miller, 1930; Villeret, 1953; Weston, 1936; Whipple and others, 1927; Yoshimura, 1931a, 1931b, 1934, 1936.

Successful irrigation requires an adequate supply of water of suitable quality, available at the proper time and place, for agricultural crop production. As an aid in conserving water, a monomolecular layer of cetyl alcohol over the entire surface of water in storage reservoirs has been advocated to reduce water evaporation (Mansfield, 1953, 1955; Rideal, 1925). However, interference in water evaporation could cause interference in oxygen atmosphere-hydrosphere interchange (Downing and Truesdale, 1956). Also, evaporation-suppressing agents might indirectly alter the iron content of stored water.

The effect of phosphorous on iron utilization by plants is reported by: Aiyar, 1948; Biddulph, 1948; Bonner and Rombyn, 1931; Brock, 1937; Brown and others, 1955; Clark, H. W., 1936; Cooper, 1939; Curry and Wilson, 1955; Davis, 1943; De and Basu, 1949; De Kock, 1955b; Dell'Aquila and Bini, 1953; Demolon and Bastisse, 1938; Einsle, 1936a, 1936b; Franco and Loomis, 1947; Goldberg, 1952; Hester and Shelton, 1947; Hopkins and Wann, 1927; Hutchinson, 1941; Iyengar, 1936; Kahler, 1956; Kapp, 1938; McDonald, 1935; Midgley, 1940; Minami and others, 1955; Ohira, 1955; Pobeguín, 1954; Sarishvili and Bagaturiya, 1938; Sideris and Young, 1956; Willis, 1936; Zicker and others, 1956.

Potassium is reported by Aiyar (1948), Eckstein and Jacob (1929), Gouny and Mazoyer (1954), Schachtschabel (1942), Scharrer and Schropp (1933), and Willis (1936) to affect the use of iron by plants.

Silicon is reported by Holl (1938), Leaf (1948), and Pobeguín (1954) to interfere in the use of iron by plants.

Aluminum is reported by Avdonin and others (1957) to interfere in the use of iron by plants.

MEASUREMENT OF MATERIALS AFFECTING IRON SOLUBILITY

Selected references pertaining to analyses of substances that alter the solubility of iron in natural water are listed in table 2.

ANALYTICAL RAMIFICATIONS OF IRON BIOCHEMISTRY

Many have found that organic matter interferes with the determination of iron in natural water (Bezel, 1948; Bode, 1932; Briggs, 1930; Clark and Sieling, 1936; Egger, 1931; Ferrey, 1935; Hallinan, 1943; Kneer, 1939; Kryukov, 1933; Marsh, 1922; Nishida, 1954;

TABLE 2.—*Humic matter, dissolved oxygen, and tannin analyses*

Substance	Material tested	Author	Remarks
Humic matter.	Soil-----	Martin and Lavollay 1950.	Permanganate oxidation of an alkali extraction.
Oxygen-----	Water-----	Faber and others, 1955.	For samples high in iron, nitrites, or organic matter (Rideal-Stewart permanganate).
	-----do-----	Gad, 1938-----	Suitable for fieldwork.
	-----do-----	Muller, 1933-----	Do.
	-----do-----	Oesting, 1934-----	Modified Van Slyke.
Tannin-----	Watercress-----	Chuprovskaya and Golota, 1954.	
	Papilionaceous plants.	Foufas, 1955----	Microchemical data are given.
		Mitchell, 1944--	Matching made with Lovibond glasses and Osborn's comparison microscope.
	<i>Spirogyra arcta</i>	Nakabayashi, 1954	Titration with K-polyvinyl alcohol sulfonate.
	<i>Potamogeton</i> spp.	Nierenstein, 1945	Method of Mitchell used.
	<i>Castanea sativa</i>	Thomas and Brossard, 1956	Color formed by gallic acid and FeCl ₃ stabilized by use of phthalic acid

Okura and Goto, 1955; Reed and Haas, 1924; Scheringa, 1931; Schnitzer and De Long, 1954a, Van Beneden, 1956a; Weston, 1936; Willcomb, 1936).

Accordingly, when organic matter is known or believed to occur in a water sample, suitable methods and techniques that take the organic matter into account should be selected for the analysis. Some methods of procedure in water sampling are discussed by Hermanowicz and Kelus (1955).

Of the methods listed in table 3, those of Bezel (1948), Briggs (1930), Clark and Sieling (1936), Hallinan (1943), Kryukov (1933), Nishida (1954), Pringsheim (1934), Scheringa (1931), Van Beneden (1956a), and Willcomb (1936) are pertinent in analysis of natural water containing organic matter.

Some iron in natural water apparently comes from atmospheric dust. Iron occurs in glacial ice and newly fallen snow (Correns, 1941; Sandstrom and Von Gegerfelt, 1945; Schenck and others, 1932; Silverman and Valenzuela, 1946; Woodcock, 1955), although the amount that could come from these sources would normally be very small. Perhaps this phenomenon is part of the reason why thorough washing of plant leaves with dilute acid is a prerequisite for valid

quantitative analyses of iron in the leaves (Jacobson, 1945; Kramer, 1957; Leonard and Stewart, 1953).

Selected references pertaining to methods for determining iron content in water, soil, and plant material are listed in table 3. They refer to likely useful analytical procedures particularly pertinent when interfering substances are present.

TABLE 3.—*Methods of analysis for iron in water, soil, and plant material*

Author	Materials tested	Method	Remarks
Aconsky and others, 1954.	Water-----	Spectrophotometric, $K_2Cr_2O_7$.	A permanent standard can be used.
Allison and Scarseth, 1942.	Soil-----	Incubate soil with sucrose under anaerobic environment.	Removes free iron oxides from soils and clays.
Am. Soc. Test. Mat., 1952.	Water-----	-----	Also tests for iron- and sulfate-reducing bacteria.
Ashizawa, 1951	do-----	Thiocyanate-----	For determining small quantities in natural water.
Astruc and Castel, 1935.	Wine-----	Fe^{++} titrated with $KMnO_4$.	Fe^{+++} reduced by Cu_2O and a small amount of H_2SO_4 .
Bashkirtseva and Yakimets, 1955.	Water-----	Titration with trilon B.	NH_4CNS is indicator.
Betremieux, 1949.	Soil-----	Glucose and NH_4 -salt extraction.	Fermentation and exchange with NH_4^+ solubilized iron in soil.
Bezel, 1948-----	Water-----	Colorimetric using thiocyanate.	Baking destroys iron humates.
Briggs, 1930-----	Medicine-----	Titration with $Na_2S_2O_3$.	Iron preparations containing organic matter.
British Standards Institute, 1956.	Water-----	-----	Industrial water testing.
Buydens and Muylle, 1952.	do-----	Thiocyanate, 0-phenanthroline, α, α' -bipyridyl.	Methods compared.
Buydens and Muylle, 1954.	do-----	Thiocyanate 2,2'-bipyridine.	Gave best reproducible results.
Carpenter and Pyle, 1934.	do-----	Fe^{++} titrated with $KMnO_4$.	
Clark and Sieling, 1936.	do-----	Colorimetric using iodohydroxyquinolinesulfonic acid.	Determination of humic iron.
Ekkert, 1931-----	do-----	Colorimetric using dimethylglyoxime.	Can be used in presence of lactate or oxalate.
Ferrey, 1935-----	do-----	Iodate titration for Fe^{++}	Method satisfactory in presence of some organic matter but not others.
Fodor and Rosenberg, 1928.	do-----	-----	Talc and kaolin adsorption of $Fe(OH)_3$.

TABLE 3.—*Methods of analysis for iron in water, soil, and plant material—Con.*

Author	Materials tested	Method	Remarks
Gad and Knetsch, 1949.	-----do-----	Similar to usual macro methods.	Analysis of very small amounts of water.
Gad and Manthey, 1950.	-----do-----	Colorimetric using KCNS, HNO ₃ and KMnO ₄ .	Method is rapid.
Goldberg and others, 1952.	Seawater-----	Microfilter followed by fixation, dehydration, and staining.	Size distribution of suspended iron particles can be studied.
Hakomori, 1931	Water-----	Colorimetric using molybdic acid, and H ₂ O ₂ , intensity of color increased by organics.	Citric, tartaric, and succinic acids interfere with determination.
Hallinan, 1943	-----do-----	Colorimetric using modified thiocyanate.	Merits are speed, ease of manipulation, and lack of interference by concentrations of organic matter.
Hetterschij, 1941	Soil-----	Extraction from soil by 1 percent citric acid or HCl at 24°C.	More iron is extracted at 24°C than at 12°C.
Hirsch and Ruter, 1926.	Water-----	Potentiometric titration.	Method determines small quantities of ferrous iron in presence of ferric iron and vice versa.
Houlihan and Farina, 1953.	-----do-----	Thiocyanate method	
Iyengar, 1937	Soil-----	Fe ⁺⁺ titrated with Ce(SO ₄) ₂ .	Diphenylaminesulfonic acid is internal indicator.
Jacobson, 1945	Plant material.	-----	Thorough washing of the leaves of plants with dilute acid is prerequisite for valid quantitative analyses of iron.
Jane, 1942	-----do-----	Microscopic technique.	Microchemical detection.
Kato, 1934	Water-----	Induced precipitation of FeS by presence of ZnS.	Due to adsorption.
Kraybill, 1930	Plant material	-----	Sample preparation.
Kretschmer, 1952	Water-----	Hellige color comparator.	Good results obtained.
Kruse, 1949	-----do-----	-----do-----	Sensitivity 0.01–0.001 mg iron per liter.
Lajoie and De Long, 1945.	Soil-----	Acid ammonium oxalate extraction.	
Lapin and Kill, 1931.	Water-----	Colorimetric using sulfosalicylic acid.	Both ferric iron and ferrous iron can be determined.
Lehmann and Reuss, 1927.	-----do-----	Colorimetric based on formation of Fe(CNS) ₃	

TABLE 3.—*Methods of analysis for iron in water, soil, and plant material—Con.*

Author	Materials tested	Method	Remarks
Leonard and Stewart, 1953.	Plant material.	-----	Citrus plant leaves should be washed before iron analysis is made.
Do -----	Water -----	Iron complexed by ethylenediamine-tetraacetic acid for removal from soil.	Precipitation of iron in acid soils prevented.
Lieber, 1953 -----	do -----	H ₂ SO ₄ , KMnO ₄ , KSCN and color comparison.	Simplified field test.
Lieffring and Buron, 1948.	do -----	Colorimetric using dimethylglyoxime.	Method is short. Limits 0-1 mg iron per liter.
Marsh, 1922 -----	Nutrient solutions.	-----	Organic iron compounds precipitate slowly from the nutrient solutions.
Meyer and Bunger, 1952.	Water -----	Pulfrich's photometer.	Determination of small amounts in a short time.
Mills, 1932 -----	Sewage -----	Microanalytical using a micro-balance.	Presence of sewage.
Morello, 1940 -----	Plant material.	Sparteine sulfate and NH ₄ CNS.	Excellent reagent for direct detection of iron.
Morgan, 1941 -----	Soil -----	0.05N CH ₃ COOH buffered at pH 4.8 with CH ₃ COONa for soil extraction.	Special tests provide indications relative to ferrous iron and ferric iron.
Morris, 1952 -----	Water -----	Tripyridyl -----	Direct determination with degree of specificity not possible by other methods.
Moss and Mellon, 1942.	do -----	Colorimetric using 2,2'-bipyridyl and 2,2',2''-terpyridyl.	
Muller, 1933 -----	do -----	-----	Field method.
New Process Rayon, Inc., 1938.	do -----	Separation of iron from copper.	Accomplished with H ₃ PO ₄ and alkali addition.
Nishida, 1954 -----	do -----	Colorimetric using KCNS.	Sample containing organic matter.
Olson, 1947 -----	Soil -----	-----	NH ₄ C ₂ H ₃ O ₂ adjusted to pH 4.8 dissolved iron from soils.
Osugi and Aoki, 1938a.	do -----	-----	At same pH more iron extracted by aqueous KCl than by HCl from soil.
Pomeroy, 1942 -----	Sewage -----	Colorimetric using NH ₄ CNS.	Sample containing more than 20 ppm iron should be diluted.

TABLE 3.—*Methods of analysis for iron in water, soil, and plant material—Con.*

Author	Materials tested	Method	Remarks
Pringsheim, 1934	Water-----	Fe ⁺⁺⁺ thiocyanate Fe ⁺⁺ K ₄ Fe(CN) ₆ total iron.	Prussian-blue test to show iron bacteria sheaths contain ferric iron.
Puri and Sarup, 1938.	Soil-----	-----	Difference in buffer solution pH makes a difference in the amount of iron extracted.
Rakestraw and others, 1936.	Seawater----	Method depends on precipitation of the iron as sulfide.	
Runehjelm, 1930	Plant material	Colorimetric using KSCN.	
Scheringa, 1931	Water-----	do-----	Sample containing pharmacopoeial iron preparations.
Schilling, 1936a	-----do-----	Fe ⁺⁺⁺ with KSCN, total iron with H ₂ O ₂ .	Iron is immediately fixed in solution with 2 ml of 1:3 H ₂ SO ₄ .
Schnitzer and De Long, 1954a.	-----do-----	2,2'-dipyridyl-----	
Scholz, 1932	Plant material	Colorimetric using NH ₄ CNS.	Determination of iron in small amounts of plant ash.
Shidlovskaya Ovchinnikova, 1953.	Water-----	Equivalent amounts of 5.5N CH ₃ COOH and 1.0N CH ₃ COONa stabilize Fe ⁺⁺ in water.	Additions of acetate buffer at pH 4.0 prevented ferrous oxidation for as long as 30 days.
Simons and others, 1953.	Seawater----	0-phenanthroline----	Sensitive and rapid.
Smith and others, 1952.	Water-----	Colorimetric using 4,7-diphenyl-1,10-phenanthroline.	Determination of 1-10γ of iron in 100 ml of water.
Steigmann, 1946	Gelatin-----	Colorimetric using P-phenylenediamine-HCl or phenyl-P-phenylenediamine-HCl.	Test is rapid.
Steinberg, 1935b	Nutrient solution.	Coprecipitation with alkaline earth as phosphate, carbonate, or hydroxide.	Method gives good results.
Svedenius, 1929	Plant material.	Precipitation with 1-nitroso-β-naphthol.	
Thompson and others, 1932.	Seawater----	Colorimetric using thiocyanate.	Interfering substances removed with H ₂ SO ₄ .
Thompson and Bremner, 1935a.	-----do-----	Colorimetric based on formation of Fe(CNS) ₃ .	
Urbach, 1934	Water-----	Step-photometric microanalysis.	
Van Beneden, 1956a.	-----do-----	Colorimetric using α,α'-dipyridyl.	Determination of humic iron.

TABLE 3.—*Methods of analysis for iron in water, soil, and plant material—Con.*

Author	Materials tested	Method	Remarks
Vanstone, 1951---	Plant material.	Selection of interval standard with proper lines suppressed.	Flame photometric analysis.
Watkins, 1954--	Water-----	Total and dissolved--	
Werescagin and others, 1931.	-----do-----	Field method-----	
Wickert and Pilz, 1950-51.	-----do-----	Detection of iron---	0-phenanthroline hydrochloride.
Wilcox, 1948-----	-----do-----	Irrigation water analysis.	
Willcomb, 1936--	-----do-----	Problem of organic matter is discussed.	Separation of iron from manganese.
Winter, 1936----	Plant material.	NH ₄ CNS is internal indicator.	Fe titrated with 0.02 N TiCl ₃ .
Yoshimura, 1936--	Water-----		Colorimetric using KSCN.
Zhivopistsev and Minina, 1954.	-----do-----	CSN- forms red precipitate solution in Me ₂ CO greatly increasing sensitivity of HCNS reaction.	Colorimetric using 2 furyl-diantipyrimylmethane.

SELECTED REFERENCES

- Abadia, A., 1956, Chlorophyll synthesis in induced iron deficiency: *Estacion Expt. Aula Dei Annales*, v. 4, p. 212-261.
- Abbott, O. D., and Ahmann, C. F., 1940, Human relationships: *Florida Soil Sci. Soc. Proc.*, v. 2, p. 109-115.
- Aconsky, L., Asami, T., and Mori, M., 1954, Permanent spectro-photometer standards for iron determination: *Am. Water Works Assoc. Jour.*, v. 46, p. 894-898.
- Adams, G. E., 1909, Weeds—their eradication and control: *Rhode Island Univ. Eng. Expt. Sta. Bull.* 133, p. 55-61.
- Adams, W. S., Leslie, A., Levin, M. H., and Luchsinger, H. W., 1950, The dermal loss of iron: *Soc. Expt. Biol. Medicine Proc.*, v. 74, p. 46-48.
- Aiyar, S. P., 1946, The place of minor elements in rice growing: *Indian Farming*, v. 7, p. 11-14.
- 1948, The effects of potassium deficiency on rice: *Indian Acad. Sci. Proc.*, v. 28B, p. 202-226.
- Aleev, B. S., and Mudretsova, K. A., 1937, The role of the phytoplankton in the nitrogen dynamics in the water of a "blossoming" pond: *Microbiology [U.S.S.R.]*, v. 6, p. 329-338.
- Alexander, C. C., and Walsh, D., 1952, New chemical agent makes possible complete control of iron chlorosis: *Agr. Chemicals*, v. 7, p. 36-38.

- Allison, L. E., and Scarseth, G. D., 1942, A biological reduction method for removing free iron oxides from soils and colloidal clays: *Am. Soc. Agronomy Jour.*, v. 34, p. 616-623.
- Allyn, W. P., 1927, The relation of lime to the absorption of iron by plants: *Indiana Acad. Sci. Proc.*, v. 37, p. 405-409.
- American Society for Testing Materials Standards, 1952, Cement, concrete, ceramics, thermal insulation, road materials, waterproofing soils: Philadelphia, Pa., Am. Soc. for Testing Materials.
- Anderson, T. G., 1940, Photoelectric colorimeter estimates of bacteria accurate: *Pennsylvania State Univ. Agr. Expt. Sta. Bull.*, v. 399, p. 14.
- Antognini, J., 1954, Iron chelates control iron chlorosis: *Agr. Chemicals*, v. 9, no. 11, p. 47-49, 131.
- Arber, A., 1920, Water plants, a study of aquatic angiosperms: Cambridge, England, Cambridge Univ. Press, 436 p.
- Arisz, W. H., and Schreuder, M. J., 1956, The path of salt transport in *Vallisneria* leaves: *Koninkl. Nederlandse Akad. Wetensch. Proc.*, ser. C, v. 59, p. 454-460.
- Arnaud, G., 1919, Treatment for chlorosis: *Rev. Viticulture*, v. 51, no. 1325, p. 325-330.
- Arndt, C. H., 1922, The growth of field corn as affected by iron and aluminum salts: *Am. Jour. Botany*, v. 9, no. 2, p. 47-71.
- Asana, R. D., 1946, Influence of nitrate, ammonium, and iron on the growth and nutrition of the rice plant: *Indian Jour. Agr. Sci.*, v. 15, p. 227-239.
- Ashizawa, T., 1951, Microanalysis of aluminum and iron: Okayama Univ. Biological Lab. Rept. no. 5, p. 76-81 [English summary]
- Askinazi, D. L., and Kheifets, D. M., 1938, Phosphates of iron and aluminum as sources of P_2O_5 for plants: *Nauchno Inst. Udobreniyam Trudy*, v. 141, p. 47-70.
- Aston, B. C., 1934, The misuse of lime with special reference to bush-sick lands: *New Zealand Jour. Agriculture*, v. 49, p. 31-36, 80-85.
- Astruc, H., and Castel, A., 1935, Determination of iron and copper in grape must and wine: *Annales Chimie anal. et Chimie appl.*, v. 17, p. 173-174.
- Atkins, W. R. G., 1923, The hydrogen ion concentration of the soil in relation to the flower color of *Hydrangea hortensis* W. and the availability of iron: *Royal Dublin Soc. Sci. Proc.*, new ser., v. 17, no. 20-24, p. 201-210.
- Austen, W., 1931, Interesting observations in the case of an iron-containing ground water with high bacterial content: *Vom Wasser*, v. 5, p. 52-56.
- Avdonin, N. S., Milovidova, E. P. Maksimova, E. D., and Frolovskaya, T. P., 1957, The influence of aluminum and manganese upon the metabolism of plants and upon the crops: *Moskovskogo Univ. Vestnik* [Moscow Univ. Bull.] Ser. Biologii, Pochvovedeniya, Geologii i. Geografii [Ser. Biology, Soil Sci., Geology, and Geography], v. 12, p. 89-97.
- Baalsrud, K., and Baalsrud, K. S., 1954, *Thiobacillus denitrificans*: *Archiv. Mikrobiologie*, v. 20, p. 34-62 [in English].
- Baas-Becking, L. G. M., Ferguson, E. J., and Kaplan, I. R., 1956, Biological processes in the estuarine environment, part 8—Iron bacteria as gradient organisms: *Koninkl. Nederlandse Akad. Wetensch. Proc.*, ser. B, v. 59, p. 398-407.
- Bass-Becking, L. G. M., and Kaplan, I. R., 1956, Biological processes in the estuarine environment, part 3—Electrochemical considerations regarding the sulfur cycle, part 4—Interpretation of observed Eh-pH relations of

- various members of the sulfur cycle: Koninkl. Nederlandse Akad. Wetensch. Proc., ser. B, v. 59, p. 85-96 [in English p. 97-108].
- Baas-Becking, L. G. M., and Mackay, M., 1956, Influence of *Enteromorpha* upon its environment: Koninkl. Nederlandse Akad. Wetensch. Proc., ser. B, v. 59, p. 109-117 [in English].
- Baas-Becking, L. G. M., and Wood, E. J. F., 1955, Biological processes in the estuarine environment, parts 1 and 2—Ecology of the sulfur cycle: Koninkl. Nederlandse Akad. Wetensch. Proc., v. 58B, p. 160-172.
- Baas-Becking, L. G. M., Wood, E. J. F., and Kaplan, I. R., 1957, Biological processes in the estuarine environment Xa-Xb. The place of the estuarine environment within the aqueous milieu: Koninkl. Nederlandse Akad. Wetensch. Proc., ser. B60, p. 88-95, 96-102.
- Baer, J. K., 1933, Disinfection of an aftergrowth in tropical drinking water: Zentralbl. gesamte Hygiene Bakteriologie u. Immunitätslehre, v. 29, p. 763-764.
- Baker, J. A., and Price, W. E., Jr., 1946, Public and industrial water supplies of the eastern coal field region of Kentucky: U.S. Geol. Survey Circ. 369, 63 p.
- Bandt, H. J., 1938, The lethal quantity of dissolved iron in fish streams: Fischerei-zeitung, v. 41, p. 593-594.
- Banfi, G., 1952, Microbes and metallic corrosion: Chimica e industria [Milan], v. 34, p. 17-21.
- Barbier, G., 1938a, Conditions and details of the fixation of colloidal ferric hydrate by soil clay: Annales Agron., v. 8, p. 34-43.
- 1938b, Ferric oxide in the clay colloids of soils: Internat. Soc. Soil Sci. Second Comm. Trans., v. 2B, p. 100-105.
- Barbier, G., and Chabannes, J., 1949, The mechanism of the adsorption of phosphates by different constituents of soil clays: Annales Agron., v. 19, p. 343-379.
- Barnette, R. M., 1921, The influence of volume upon the rate of change in contact with plant roots and the effect of this change upon iron availability: New Jersey State Agr. Expt. Sta. Rept., p. 345-347.
- Barnette, R. M., and Shive, J. W., 1923, The influence of solution volume upon plant growth in relation to reaction change and iron availability in culture solutions: Soil Sci., v. 15, no. 5, p. 413-425.
- Bashkirtseva, A. A., and Yakimets, E. M., 1955, The use of Trilon B for the determination of iron in natural waters: Zavodskaya Lab., v. 21, p. 533-535.
- Baudenistel, R., 1957, Chelated iron corrects chlorosis of gardenias: New York State Flower Growers Bull. 143, p. 2-3.
- Baylis, J. R., 1925, Natural water corrosion and hydrogen-ion concentration: Chem. Metall. Eng., v. 32, p. 874-875.
- Bear, F. E., 1954, Progress report on research with particular reference to New Jersey soils: Jour. Agr. Food Chemistry, v. 2, p. 244-251.
- Beaumont, A. B., and Holland, E. B., 1935, Absorption by food plants of chemical elements of importance in human nutrition: Massachusetts Agr. Expt. Sta. Bull. 315, p. 14-15 [Ann. rept., 1934].
- Beger, H., 1935, *Leptothrix echinata*, a new iron bacterium which can precipitate manganese: Zentralbl. Bakteriologie Parasitenkunde, Abt. 2, v. 92, p. 401-406.
- 1937, The iron bacteria in water systems and their practical significance: Gas-u. Wasserfach, v. 80, p. 886-889, 908-911.

- Beger, H., and Haase, L. W., 1941, Iron and manganese disturbances in a water works; causes and treatment: Chem. Zentralbl. (Leipzig, Berlin) 1942, v. 1, p. 20-49.
- Belcher, R., Gibbons, D., and West, T. S., 1955, Effect of ethylenediaminetetraacetic acid on the ferrous/ferric and cuprous/cupric oxidation-reduction systems: Anal. Chimica Acta, v. 12, p. 107-114.
- Bennett, J. P., 1928, Plant pathology: California Agr. Expt. Sta. Rept., p. 40-44, 89-93, 96, 97.
- 1945, Iron in leaves: Soil Sci., v. 60, p. 91-105.
- Bertrand, D., and De Wolf, A., 1956, Importance of iron as a trace element in the utilization of citric acid by *Aspergillus niger*: Acad. sci. [Paris] Comptes rendus, v. 243, p. 1924-1926.
- Betremieux, R., 1949, Fermentation of organic matter and chemical dynamics of the industrial soil: Acad. sci. [Paris] Comptes rendus, v. 228, p. 1749-1751.
- 1951, Evolution of iron and manganese in the soil: Inst. Natl. recherche agron. [France] Annales, ser. A, v. 2, p. 193-295.
- 1954, Effect of organic matter fermentation on the status of mineral elements in undisturbed soils: Acad. sci. [Paris] Comptes rendus, v. 238, p. 2257-2258.
- Beutelspacher, H., 1955, Reciprocal effect between inorganic and organic colloids of soil: Zeitschr. Pflanzenernährung Düngung Bodenkunde, v. 69, p. 108-115.
- Bezel, L. I., 1948, Determination of iron in colored waters: Zavodskaya Lab., v. 14, p. 799-800.
- Bezier, D., 1943, Oxidation-reduction potential of ferric ion solutions with very low concentrations of ferrous ions: Acad. sci. [Paris] Comptes rendus, v. 216, p. 798-800.
- Bhaskaran, T. R., and Pillai, S. C., 1937, Decomposition of cane molasses in the soil and its bearing on soil fertility: Soc. Biol. Chemists [India] Proc., v. 2, p. 33-34.
- Biddulph, O., 1948, Interrelations between iron and phosphorous in plant nutrition: Oregon State Chap. Phi Kappa Phi, Corvallis, Biology Colloquium, Nutrition, p. 24-27.
- Black, W. A. P., and Mitchell, R. L., 1952, Trace elements in the common brown algae and in sea water: Marine Biol. Assoc. United Kingdom Jour., v. 30, p. 575-584.
- Blagovidov, N. L., Rabinovich, V. A., and Sell-Bekman, I. Y., 1957, The changes of the oxidation-reduction potential in the profile of several soils in the Leningrad province: Pochvovedenie, 6, p. 81-85.
- Bloomfield, C., 1952, Translocation of iron in Podzol formation: Nature, v. 170, p. 540.
- 1957, The solution-reduction of ferric oxide by aqueous leaf extracts: The role of certain constituents of the extracts: Internat. Cong. Soil Sci., 6th Trans. [Paris], v. B, p. 427-432.
- Blum, J. L., 1953, The ecology of algae growing in the Saline River, Michigan, with special reference to water pollution: Unpublished doctor of philosophy dissertation, Michigan Univ., 176 p.
- 1956, The ecology of river algae: Bot. Rev., v. 22, p. 291-341.
- Blunno, M. L., 1911, On the proportion of iron sulphate used against white rot of grapevines: Australian Assoc. Adv. Sci. Rept., v. 13, p. 562-566.

- Bochkarev, P. F., and Yanko, A. P., 1953a, Hydrochemical characteristics of Lake Altarik: Akad. Nauk SSSR, Inst. obshch. neorgan. khimii, Izv. sektora fiziko-khim. analiza, v. 1, no. 1/2, p. 225-228.
- 1953b, Hydrochemical characteristics of the Baushevskii Pond: Irkutsk Univ. Nauchno Issledovatel'skogo Inst., Izv. fiziko-khim. v. 1, no. 1/2, p. 229-237.
- Bode, G., 1932, Unionized iron in subsoil water: Wochenschr. Brauerei, v. 49, p. 321-322.
- Boiret, H., and Paturel, G., 1892, The use of sulphate of iron in agriculture: Annales Agron., v. 18, p. 417-440.
- Boischot, P., and Durroux, M., 1949, Fixation of iron and manganese in soils containing lime: Acad. sci. [Paris] Comptes rendus, v. 229, p. 380-381.
- 1950, Movement of iron and manganese in calcareous soils: Annales Agron., v. 1, p. 551-554.
- Bollmann, A., and Schwanitz, F., 1957, The iron content of plants from an iron rich habitat: Zeitschr. Botanik, v. 45, p. 39-42.
- Bonner, J. F., 1950, Plant biochemistry: New York, Academic Press, 537 p.
- Bonner, W. D., and Rombyn, H., Jr., 1931, Oxidation of ferrous iron by iodine in the presence of phosphate and the nonexistence of a ferric phosphate complex: Indus. Eng. Chemistry, Anal. Ed., v. 3, p. 85-87.
- Bortels, H., 1927, The significance of iron, zinc, and copper for microorganisms: Zeitschr. Biochemie, v. 182, p. 301-358.
- 1953, Supplemental trace elements concerned with vibrios and cytophages: Zentralbl. Bakteriologie Parasitenkunde Abt. 1 Orig. v. 160, p. 213-216.
- Bould, C., 1957, Iron chelates in plant nutrition: Jour. Sci. Food Agriculture, v. 8, [supp. issue], p. S1-S5.
- Bowen, W. J., 1940, Effects of vanadium, copper, manganese, and iron on the synthesis of protoplasm by *Chilomonas paramecium*: Biol. Bull., v. 79, p. 114-130.
- Bowman, H. H. M., 1956, Salinity data on marine and inland waters and plant distribution: Ohio Jour. Sci., v. 56, p. 101-106.
- Braley, S. A., 1957, Evaluation of mine-water drainage: Am. Inst. Mining Metall. Petroleum Engineers Trans. 208, Tech. Pub. 4414-F (in Mining Eng. 9, no. 1, p. 76-78).
- Bray, W. C., and Hershey, A. V., 1934, The hydrolysis of ferric ion—The standard potential of the ferric-ferrous electrode at 25° C—The equilibrium $\text{Fe}^{+++} + \text{Cl} = \text{FeCl}^{++}$: Am. Chem. Soc. Jour., v. 56, p. 1889-1893.
- Brewer, P. H., and Carr, R. H., 1927, Fertility of a soil as related to the forms of its iron and manganese: Soil Sci., v. 23, no. 3, p. 165-173.
- Briggs, W. P., 1930, Assays of some official iron preparations containing organic matter: Am. Pharmaceut. Assoc. Jour., v. 19, p. 1191-1198.
- British Standards Institute, 1956, Methods of testing water used in industry: British Standards Inst., London, S. W. 1, British Standard 2690, 150 p.
- Broadbent, F. E., 1954, Modification in chemical properties of straw during decomposition: Soil Sci. Soc. America Proc., v. 18, p. 165-169.
- Brock, J. F., 1937, The effect of large doses of iron on the absorption of phosphorus: Clinical Sci., v. 3, p. 37-45.
- Brown, H. J., and Gibbons, N. E., 1955, Effect of magnesium, potassium, and iron on the growth and morphology of red halophilic bacteria: Canadian Jour. Microbiology, v. 1, p. 486-494.

- Brown, J. C., and Holmes, R. S., 1955, Iron, the limiting element in chlorosis, part 1—Availability and utilization of iron dependent upon nutrition and plant species: *Plant Physiology*, v. 30, p. 451-457.
- Brown, J. C., Holmes, R. S., Shapiro, R. E., and Specht, A. W., 1955, Effects of phosphorus and copper salts on iron chlorosis of rice in flooded and non-flooded soil and the associated enzymic activity: *Soil Sci.*, v. 79, p. 363-372.
- Brown, L. A., 1934, Oxidation-reduction potentials in soils, part 1—Principles and electrometric determination: *Soil Sci.*, v. 37, p. 65-76.
- Brown, R., and Possingham, J. V., 1957, Iron deficiency and the growth of pea roots: *Royal Soc. [London] Proc.*, v. B 147, p. 145-166.
- Brozek, Bohumil, 1949, Changes in the chemical composition of mineral waters during storage: *Lekarske Listy*, v. 4, p. 386-391.
- Bruevich, S. V., and Vinogradova, E. G., 1946, General features of sedimentation in the Caspian Sea (according to the distribution of carbonates, iron, manganese, and phosphorus in sea deposits): *Acad. sci. [U.S.S.R.] Comptes rendus*, v. 52, p. 789-792.
- Brussoff, A., 1918, A calcium and iron metabolizing bacillus: *Centralbl. Bakteriologie Parasitenkunde*, Abt. 2, v. 48, p. 193-210.
- Bryner, L. C., Beck, J. V., Davis, D. B., and Wilson, D. G., 1954, Microorganisms in leaching sulfide minerals: *Indus. Eng. Chemistry*, v. 46, p. 2587-2592.
- Burk, D., Lineweaver, H., and Horner, K. C., 1932, Iron in relation to the stimulation of growth by humic acid: *Soil Sci.*, v. 33, p. 413-453.
- Burk, D., Lineweaver, H., Horner, C. K., and Allison, F. E., 1931, The relation between iron, humic acid, and organic matter in the nutrition and stimulation of plant growth: *Science*, v. 74, p. 522-524.
- Butcher, R. W., 1933, Studies on the ecology of rivers, part 1—On the distribution of macrophytic vegetation in the rivers of Britain: *Jour. Ecology*, v. 21, p. 58-91.
- Butlin, K. R., Vernon, W. H. J., and Whiskin, L. C., 1952, Investigations on underground corrosion: *Water and Water Eng.*, January 1952, p. 15-18.
- Buydens, R., and Muylle, R., 1952, Determination of iron in water: *Centre Belge etude doc. eaux Bull. [Liège]* 18, p. 241-245.
- 1954, Determination of iron in water: *Centre Belge etude doc. eaux Bull. [Liège]* 23, p. 52-55.
- Byram, A. T., 1938, Chemical water analyses and their interpretation: *Eng. Contract Rec.*, v. 51, no. 12, p. 42-44.
- Calgon, Inc., and Albright and Wilson, Ltd., 1944, Preventing the precipitation of iron dissolved in water: *British Patent No. 560,905*.
- California State Water Pollution Control Board, 1952, Water quality criteria: *California State Water Pollution Control Board Pub. 3*, p. 276-278.
- Calise, V. J., and Dietz, R. F., 1955, Iron and manganese removal from municipal and industrial water supplies: *Graver Water Conditioning Co. Tech. Repr. T-144*, 8 p.
- Cameron, E. J., 1934, "Blackbeets," a problem involving stimulation of bacterial growth by iron: *Jour. Bacteriology*, v. 27, p. 60-61.
- Carnes, G., 1954, Has your plant had its iron lately? Try a chelating agent: *New Jersey Agriculture*, v. 36, no. 4, p. 8-9, 11.
- Carpenter, L. V., and Pyle, G. R., 1934, Lime demand of acid water: *Pennsylvania Water Works Assoc. Jour.*, v. 6, p. 47-48.
- Cataldi, M. S., 1937, Iron bacteria in the drinking water of Buenos Aires: *Bol. Obras Sanitarias Nacion [Buenos Aires]*, v. 1, p. 464-478.

- Cervenka, R., 1955, Influence of copper and of its compounds on algae: *Gaz, Woda i Tech. Sanitarna*, v. 29, p. 66-69.
- Chaberek, S., Jr., and Bersworth, F. C., 1953, New chelating agents for trivalent iron: *Science*, v. 118, p. 280.
- Chernovskaya, E. N., 1954, Oxygen content of White Sea water: *Akad. Nauk. SSSR Doklady*, v. 97, p. 911-913.
- Chittenden, R. H., 1896, The amount of iron in ordinary dietaries: *New York, Dietetic and Hygienic Gaz.*, v. 12, no. 1, p. 24-26.
- Cholodnii, N., 1928, So-called iron organisms and the Naumann method for their investigation: *Deutsche Bot. Gesell. Ber.*, v. 46, p. 317-323.
- Chuprovskaya, S. V., and Golota, G. M., 1954, Phytochemical studies of water-cress: *L'vov, Med. Inst. Rabot Nauchnoe Studencheskoe Obshch.*, Sbornik 2, p. 61-63.
- Clark, F., Nearpass, D. C., and Specht, A. W., 1957, Influence of organic additions and flooding on iron and manganese uptake by rice: *Agronomy*, v. 49, p. 586-589.
- Clark, H. W., 1936, Occurrence of iron in ground water and early experiments upon removal: *New England Water Works Assoc. Jour.*, v. 50, p. 229-230.
- Clark, N. A., 1936, Green plants and the reactions of iron: *Iowa Acad. Sci. Proc.*, v. 43, p. 185-189.
- Clark, N. A., and Sieling, D. G., 1936, Determination of iron in humates—Use of Iodo-hydroxyquinolinesulfonic acid: *Indus. Eng. Chemistry, Anal. Ed.*, v. 8, p. 256-257.
- Clarke, F. W., 1924, The composition of the river and lake waters of the United States: *U.S. Geol. Survey Prof. Paper* 135.
- Cohn, M. L., Kovitz, C., Oda, V., and Middlebrook, G., 1954, Isoniazid and tubercle bacilli, part 2—The growth requirements, catalase activities, and pathogenic properties of isoniazid-resistant mutants: *Am. Rev. Tuberculosis*, v. 70, p. 641-664.
- Coldwell, B. B., and De Long, W. A., 1950, Studies on the composition of deciduous forest tree leaves before and after partial decomposition: *Sci. Agriculture*, v. 30, p. 456-466.
- Cooper, L. H. N., 1935, Iron in the sea and in marine plankton: *Royal Soc. [London] Proc.*, v. B 118, p. 419-438.
- 1939, Phosphorus, nitrogen, iron, and manganese in marine zooplankton: *Marine Biol. Assoc. United Kingdom Jour.* v. 23, p. 387-390.
- 1948, Some chemical considerations on the distribution of iron in sea water: *Marine Biol. Assoc. United Kingdom Jour.*, v. 27, p. 314-321.
- Copeland, O. L., Jr., 1957, Influence of moisture and compaction on soil oxidation reduction potentials: *Soil Sci. Soc. America Proc.*, v. 21, p. 269-71.
- Correns, C. W., 1941, Geochemistry of manganese and iron: *Akad. Wiss, Göttlingen Nachr., Math.-Phys. Kl.*, no. 5, p. 219-230.
- Coste, F., 1896, The role of sulphate of iron: *Prog. Agricole Viticole*, v. 26, no. 42, p. 434-440; no. 43, p. 463-467; no. 44, p. 496-504.
- Coupin, H., 1898, On the poisonous effect of the salts of copper on the higher plants: *Acad. sci. [Paris] Comptes rendus*, v. 127, no. 10, p. 400-401.
- Crabill, M. P., 1956, Biologic infestation at Indianapolis: *Am. Water Works Assoc. Jour.*, v. 48, p. 269-274.
- Crafts, A. S., 1954, Composition of the sap of xylem and phloem and its relation to the nutrition of the plant: *Cong. Internat. Bot.*, 8th, Paris, Colloques analyse plantes et problemes engrais minéraux, p. 18-21 [in English].

- Crawford, R. F., 1939, The causes and control of chlorosis in New Mexico: New Mexico Agr. Expt. Sta. Bull. 264, 12 p.
- Curini-Galletti, A., 1937, The yellowing of wheat seedlings under vine rows: Riv. Patologia Vegetale, v. 27, p. 149-160.
- Curry, J. J., and Wilson, S. L., 1955, Effect of sewage-borne phosphorus on algae: Sewage Indus. Wastes, v. 27, p. 1262-1266.
- Dalle, R. A. D., 1946, Function of biogeneous elements and soil exhaustion: Italian Agriculture, v. 83, p. 744-746.
- Datsko, V. G., 1955, The contents of organic substances in the Azov Sea before the regulation of the discharge of the Don River: Hidrokhim. Materialy, v. 23, p. 3-10.
- Davis, F. L., 1943, The role of iron and aluminum in the retention of phosphates by soils as indicated by the solubility of phosphorus: Soil Sci. Soc. America Proc., v. 8, p. 167-170 [1944].
- Davis, G. K., and Loosli, J. K., 1954, Mineral metabolism: Ann. Rev. Biochemistry, v. 23, p. 459-480.
- De, H. N., and Basu, K. P., 1949, Mutual influence of minerals in metabolism: Indian Jour. Med. Research, v. 37, p. 213-231.
- De Boissezon, P., 1933, Utilization of proteins and iron of vegetable origin in the maturation of eggs of *Culex pipiens*: Soc. Biologie [France] Comptes rendus, v. 114, p. 487-489.
- Deevey, E. S., Jr., 1941, Limnological studies in Connecticut, part 7—The quantity and composition of the bottom fauna of 36 Connecticut and New York lakes: Ecol. Mon., v. 11, p. 413-455.
- De Gruchy, J. H. B., 1938, A preliminary study of the larger aquatic plants of Oklahoma with special reference to their value in fish culture: Oklahoma Agr. Mech. Coll. Agr. Expt. Sta., Stillwater, Okla., Tech. Bull. 4.
- De Jesus, P. I., 1937, Physicochemical factors in anopheline ecology, part 2—Studies on turbidity, chloride, and iron: Philippine Jour. Sci., v. 62, p. 125-142.
- De Kock, P. C., 1955a, Iron nutrition of plants at high pH: Soil Sci., v. 79, p. 167-174.
- 1955b, Influence of humic acids on plant growth: Science, v. 121, p. 473-474.
- 1955c, Chlorosis and the iron nutrition of plants: Cong. Internat. Biochimie 3d [Brussels], p. 102.
- Delecourt, J., 1947, The presence of iron in waters from certain wells: Technique l'eau [Brussels], p. 11-12.
- Dell'Aquila, D., and Bini, G., 1953, Influence of calcium on iron absorption in man: Riforma Medica, v. 67, p. 1073-1078.
- De Long, W. A., and Schnitzer, M., 1955, Investigations on the mobilization and transport of iron in forested soils, part 1—The capacities of leaf extracts and leachates to react with iron: Soil Sci. Soc. America Proc., v. 19, no. 3, p. 360-363.
- Dementiev, K. I., 1941, Ferrobacteria in Old Russian mineral springs and lakes: Balneological observation sta. of the resort "Storaja Rossia," Leningrad Province [English summary].
- Demolon, A., and Bastisse, E., 1938, Migration of iron in soils: Cong. Chim. Indus. Comptes rendus, 18th Cong., Nancy, p. 154-157.
- Densch, A., and Hunnius, T., 1924, Experiments with copper sulfate: Zeitschr. Pflanzenernährung Düngung, v. 3, no. 6, p. 369-386.

- Derby, R. L., 1956, Chlorination of deep reservoirs for taste and odor control: *Am. Water Works Assoc. Jour.*, v. 48, p. 775-780.
- De Wilkoszewski, B., 1917, An agronomic study of several fertilizing or anti-cryptogamic compounds used in agriculture: *Archives Sci. Phys. Nat.* [Geneva], ser. 4, v. 44, p. 165-180.
- Dickey, R. D., 1942, Iron deficiency of tung in Florida: *Assoc. Southern Agr. Workers, Proc. Ann. Convention*, v. 43, p. 173-174.
- Dienert, F., 1938, Iron and manganese in water: *Technique sanitaire municipale*, v. 33, p. 105-116.
- Dion, H. G., 1944, Iron oxide removal from clays and its influence on base-exchange properties and X-ray diffraction patterns of the clays: *Soil Sci.*, v. 58, p. 411-424.
- Downing, A. L., and Truesdale, G. A., 1956, Aeration in aquaria: *Zoologica*, v. 41, p. 129-143.
- Doyne, H. C., and Morison, C. G. T., 1926, The absorption of iron by soils: *Soil Sci.*, 22, no. 3, p. 163-173.
- Dubois, L., 1955, Water hyacinth in the Belgian Congo: *Bull. Agr. Congo Belge*, v. 46, p. 893-900.
- Duchaufour, P., 1953a, Note on iron solubility by organic matter in certain soils: *Assoc. Française Étude Sci. Bull.*, v. 17, no. 6, p. 483.
- 1953b, Applications of pedology in the tropics: *Soc. Sci. Bull.*, Nancy, v. 21, p. 19-22.
- Duchon, K., and Miller, L. B., 1948, Effect of chemical agents on iron bacteria: *Paper Trade Jour.*, v. 126, no. 4, p. 47-58.
- Dupont, P., 1954, Vegetation of the marshes at the mouth of the Vilaine River: *Soc. Sci. Bretagne Bull.*, v. 29, p. 65-104 [1955].
- Dusserre, C., 1900, Destruction of weeds in fields of cereals: *Annuaire agr. Suisse*, v. 1, no. 9, p. 331-337.
- Dyakonova, K. V., 1957, Soil as a source of carbonic acid (carbon dioxide) for plants in irrigated and non-irrigated ciscaucasian chernozems: *Pochvovedenie*, 10, p. 86-92.
- Eames, A., and MacDaniels, L. H., 1925, An introduction to plant anatomy: New York, McGraw-Hill, p. 309-314.
- Echave, D., 1937, Copper, a biogenetic element in vegetables. Its determination in some plant species in Argentina: *Rev. Farmaceutica* [Buenos Aires], v. 79, p. 81-84.
- Eckstein, O., and Jacob, A., 1929, The potassium oxide-iron antagonism in the plants as basis of a method for determining the potassium oxide requirements of soils: *Zeitschr. Pflanzenernährung, Düngung Bodenkunde*, v. 14A, p. 205-220.
- Edelstein, F., and Von Csonka, F., 1912, The iron content of cow's milk: *Biochem. Zeitschr.*, v. 38, no. 1-2, p. 14-22.
- Edgerton, L. J., 1942, Use of metaphosphate in nutrient solutions: *Am. Soc. Horticultural Sci. Proc.*, v. 41, p. 237-239.
- Edmonson, W. T., Anderson, G. C., and Peterson, U. R., 1956, Artificial eutrophication of Lake Washington: *Limnology and Oceanography*, v. 1, no. 1, p. 47-53.
- Egger, F., 1931, Iron and manganese-containing corrosive waters: *Vom Wasser*, v. 5, p. 46-52.
- Eggleton, P., 1931, Recent advances in science: *Biochemistry: Sci. Prog.*, v. 26, p. 16-21.

- Einsele, W., 1936a, Relation of iron circulation to phosphate circulation in eutrophic lakes: *Archiv. Hydrobiologie*, v. 29, p. 664-685.
- 1936b, The significance of iron, pH, and decomposition in the utilization of phosphate (as fertilizer in fish ponds): *Fischerei-zeitung*, v. 39, p. 774-776.
- 1940a, A search for a theory of the dynamics of the manganese and iron stratification in an eutrophic lake: *Naturwissenschaften*, v. 28, p. 257-264.
- 1940b, Theory of the dynamics of the manganese and iron layers in eutrophic seas: *Naturwissenschaften*, v. 28, p. 280-285.
- Ekkert, L., 1931, A test for the ferrous ion: *Magyar Gyogyszeresztudomanyi Tarsasag Ertesitoje*, v. 7, p. 231-232.
- Ellis, D., 1919, *Iron bacteria*: London, Methuen and Co., Ltd., 179 p.
- 1921, Iron bacteria in relation to the incrustation of pipes: *Engineering*, v. 112, p. 457-458.
- 1925, Iron incrustation in water pipes—The responsibility of iron algae: *Water and Water Eng.*, v. 27, p. 352-354.
- Elvehjem, C. A., 1931, The role of iron and copper in the growth and metabolism of yeast: *Jour. Biol. Chem.*, v. 90, p. 111-132.
- 1935, The biological significance of copper and its relation to iron metabolism: *Physiol. Rev.*, v. 15, p. 471-507.
- Elvehjem, C. A., Herrin, R. C., and Hart, E. B., 1927, Iron in nutrition, part 3—The effects of diet on the iron content of milk: *Jour. Biol. Chemistry*, v. 71, no. 2, p. 255-262.
- Emmons, W. H., Thiel, G. A., Stauffer, C. R., and Allison, I. S., 1955, *Geology, principles and processes*: 4th ed. New York, McGraw-Hill, 638 p.
- Enari, T. M., 1955, Effect of cobalt and iron on riboflavine production by *Candida guilliermondia*: *Acta Chemica Scandinavica*, v. 9, p. 1726-1727 [in English].
- English, M. P., and Barnard, N. H., 1955, The effect of tracemetal deficiency on some *Trichophyton* strains: *British Mycological Soc. Trans.*, v. 38, p. 78-82.
- Eristavi, D. I., 1953, Oxidation and reduction in water chemistry: *Politekhnikeskogo Inst. [Budapest] Trudy*, v. 28, p. 31-42.
- 1955, Oxidation and reduction in hydrochemistry: *Gidrokhim. Materialy*, v. 24, p. 27-29.
- Erkama, J., 1949, Effect of copper on the iron uptake of plants: *Acta Chemica Scandinavica*, v. 3, p. 850-857 [in English].
- Escudero, A., Herraliz, M. L., and Senra, R., 1944, Study of the nutritive value of cress: *Assoc. Argentina Dietologia Rev.*, 2, 5, p. 54-56.
- Evenson, M. A., and Gerhardt, P., 1955, Nutrition of *Brucellae*; utilization of iron, magnesium, and manganese for growth: *Soc. Expt. Biology Medicine Proc.*, v. 89, p. 678-680.
- Faber, H. A., Hatfield, W. D., McCrady, M. H., and Kline, E. K., 1955, *Standard methods for the examination of water, sewage, and industrial wastes*: 10th ed. New York, Am. Public Health Assoc., 522 p.
- Fabrin, A. O., 1948, Selection of deep well pumps: *Am. Water Works Assoc. Jour.*, v. 40, p. 415-422.
- Falci, M. M., 1935, The antagonistic action of an iron mineral water to sparteine, strychnine, and morphine: *Rassegna Terapia Patologia Clinica*, v. 7, p. 1-8.
- Fassett, N. C., 1940, *A manual of aquatic plants*: New York, McGraw-Hill, 382 p.
- Fattinger, D., 1950, The influence of other ions on the fungicidal activity of the copper ion: *Pflanzenschutz Ber.*, v. 4, p. 33-46.

- Feeney, R. E., Mueller, J. H., and Miller, P. A., 1943, Growth requirements of *Clostridium tetani*, part 2—Factors exhausted by growth of the organism: Jour. Bacteriology, v. 46, p. 559-562.
- Ferrey, G. J. W., 1935, Determination of ferrous iron in presence of organic matter by Heisig's method: Quart. Jour. Pharmacy Pharmacology, v. 8, p. 344-350.
- Fischer, E. K., 1950, Colloidal dispersions: New York, Wiley and Sons, 387 p.
- Fly, C. L., 1935, Organic iron and hydrogen-ion concentration as associated factors affecting the rate of reproduction of *Lemna major*: Oklahoma Acad. Sci. Proc., v. 15, p. 77-80.
- Fodor, A., and Rosenberg, A., 1928, The adsorption of iron by kaolin and talc from solutions of ferric hydroxide and chloride: Kolloid-Zeitschr., v. 46, p. 91-95.
- Fogg, G. E., 1952, The metabolism of algae: New York, Wiley and Sons, p. 114.
- Follett-Smith, R. R., 1938, Flood-fallowing of sugar cane soils: Internat. Soc. Sugar-Cane Technology Proc., v. 6, p. 840-842.
- Follett-Smith, R. R., and Robinson, L. A., 1936, Flood fallowing of sugar cane soils: Internat. Soc. Sugar-Cane Technology Proc., v. 7, p. 227-230.
- Forman, Jonathan, and Frink, O. E., 1950, Water and Man; a study in ecology: Columbus, Ohio, Friends of the Land, p. 284-295.
- Forster, W. A., 1953, Some effects of metals in excess on crop plants grown in soil culture, part 6—Effects of cobalt and copper on mineral status of oat: Agr. Hort. Research Sta., Long Ashton, Bristol, p. 214-218.
- Foufas, C. A., 1955, Investigation of tannic substances in papilionaceous plants: Akad. Athenon Praktika, v. 30, p. 302-306.
- Francis-Boeuf, C., 1948, The possibility of a physiology of marine sediments: Museum Histoire Nat. [Marseille] Bull., v. 8, no. 1, p. 37-43.
- Franco, C. M., and Loomis, W. E., 1947, The absorption of phosphorus and iron from nutrient solutions: Plant Physiology, v. 22, p. 627-634.
- Free, A. H., 1940, Non-identity of gray hair production by mineral deficiency and vitamin deficiency: Soc. Expt. Biology Medicine Proc., v. 44, p. 371-373.
- Frisbie, R. I., 1953, The biochemistry of iron: Gaceta Medica Mexico, v. 83, p. 389-400.
- Fry, E. J., 1924, A suggested explanation of the mechanical action of lithophytic lichens on rocks (shale): Annals Botany, v. 38, p. 175-196.
- Fujihara, T., 1925, Carbon dioxide in corrosion of iron: Chem. Metall. Eng., v. 32, p. 810-811.
- Fujii, Y., and Tanaka, N., 1955, Absorption of oxygen by seedlings of rice plants of various varieties: Crop Sci. Soc. Japan Proc. [Tokyo], v. 24, p. 95-96.
- Gad, G., 1938, A simple colorimetric method for determination of dissolved oxygen in water: Gas- u. Wasserfach, v. 81, p. 59-60.
- Gad, G., and Knetsch, M., 1949, Chemical analysis of very small amounts of water: Gesundheits-Ingenieur, v. 70, p. 259-261.
- Gad, G., and Manthey, M., 1950, Determination of total iron in water: Gesundheits-Ingenieur, v. 71, p. 27-29.
- Galtsoff, P. S., 1934, The biochemistry of the invertebrates of the sea: Ecol. Mon., v. 4, p. 481-490.
- Gantimurov, I. I., 1952, Solontes and marsh processes in the Om basin: Pochvovedenie 1007-11 (R), Soils and Fert. 16, no. 2, p. 111.
- Garibaldi, J. A., and Neilands, J. B., 1955, Isolation and properties of ferri-chrome A: Am. Chem. Soc. Jour., v. 77, p. 2429-2430.

- Gausman, H. W., and Struchtemeyer, R. A., 1957, Effects of fertilizers and restricted aeration on the subterranean morphology of the potato plant: *Am. Potato Jour.*, v. 34, p. 285-286.
- Gayral, P., 1954, Phytolimnological investigations in Morocco: *Soc. Sci. Nat. Phys. Maroc Travaux Inst. Sci. Cherifien, Ser. Bot.* 4, 306 p.
- Gayral, P., and Panouse, J. B., 1954, Physical and biological researches on Lake Aguelmame Azigza; *Soc. Sci. Nat. Phys. Maroc Bull.*, v. 34, p. 135-159.
- Gellerman, V. M., 1950, Effect of decomposition products of organic substances on root nutrition of plants: *Akad. Nauk SSSR Doklady*, v. 73, p. 593-596.
- Gerb, L., 1956, Chemical processes in ground-water wells: *Gas- u. Wasserfach*, v. 97, p. 607-610.
- Gerloff, G. C., and Skoog, F., 1957, Availability of iron and manganese in southern Wisconsin lakes for the growth of *Microsystis-aeruginosa*: *Ecology*, v. 38, p. 551-556.
- Gessner, F., 1956, Water chemistry and water vegetation: *Urania*, v. 19, p. 328-337.
- Gile, P. L., and Carrero, J. O., 1914, Assimilation of colloidal iron by rice: *U.S. Dept. Agriculture Jour. Agr. Research*, v. 3, no. 3, p. 205-210.
- 1916a, Immobility of iron in the plant: *U.S. Dept. Agriculture Jour. Agr. Research*, v. 7, no. 2, p. 83-87.
- 1916b, Assimilation of iron by rice from certain nutrient solutions: *U.S. Dept. Agriculture Jour. Agr. Research*, v. 7, no. 12, p. 503-528.
- Gleen, H., 1950, Biological oxidation of iron in soil: *Nature*, v. 166, p. 871-872.
- Glenk, H. O., 1954, Seasonal changes in the phytoplankton of the Dechsendorf Pond near Erlangen: *Phys. Med. Sozietat Erlangen Sitzungsber.*, v. 77, p. 58-157 [1956].
- Glicklhorn, J., 1920, Iron organisms, part 1: *Akad. Wiss. Wien Sitzungsber., Abt. 1*, v. 129, p. 187-213.
- Gloyna, E. F., Hermann, E. R., and Drynan, W. R., 1955, Oxidation ponds—waste treatment studies, radioisotope uptake and algae concentration: *U.S. Atomic Energy Comm., AECU-3113*, 90 p.
- Goldberg, E. D., 1952, Iron assimilation by marine diatoms: *Biol. Bull.*, v. 102, p. 243-248.
- Goldberg, E. D., Baker, M., and Fox, D. L., 1952, Microfiltration in oceanographic research, part 1—Marine sampling with the molecular filter: *Jour. Marine Research (Sears Foundation)*, v. 11, p. 194-204.
- Goldschmidt, V. M., 1954, *Geochemistry*: Oxford, England, Clarendon Press, p. 662-667.
- Gollmick, F., 1936, The influence of zinc, iron, copper, and their combinations on the growth of *Aspergillus niger*: *Zentralbl. Bakteriologie Parasitenkunde, Abt. 2*, v. 93, p. 421-442.
- Gorbunov, K. V., 1957, Relations between aerobic and anaerobic processes in meadow soils and water used for flooding: *Mikrobiologiya*, v. 26, p. 210-217.
- Gorham, A. V., and Gorham, E., 1955, Iron, manganese, ash, and nitrogen in some plants from salt marsh and shingle habitats: *Annals Botany [London]*, v. 19, p. 571-577.
- Gortner, R. A., 1934, Lake vegetation as a possible source of forage: *Science*, v. 80, p. 531-533.
- Goryunova, S. V., and Nasonova, M. V., 1955, Amount and character of phytoplankton from Lake Beloe Ozero in different seasons: *Mikrobiologiya*, v. 24, p. 435-443.

- Gouny, P., Mazoyer, R., 1954, Relations between mineral nutrition and pathological symptoms of lime-induced chlorosis: Soils and Fertilizers [Harpden, England], v. 17, p. 51.
- Graftiau, J., 1900, Combating weeds by means of chemical agents: Ingenieur Agricole Gembloux, v. 10, no. 6, p. 413-428.
- Granick, S., 1954, Iron metabolism: New York Acad. Medicine Bull., v. 30, p. 81-105.
- Greaves, J. E., 1933, Some factors influencing nitrogen fixation: Soil Sci., v. 36, p. 267-280.
- Greer, E. N., Pringle, W. J. S., and Kent, N. L., 1952, Composition of British grown winter wheat, part 2—Iron and manganese contents: Sci. Food Agriculture Jour., v. 3, p. 16-18.
- Griffin, A. E., 1937, The role of chlorine in water purification: Ann. Conf. Water Purification in West Virginia, 11th, Proc.: West Virginia Univ. Eng. Expt. Sta., Tech. Bull. 9, p. 18-23.
- Griffith, R. E., 1955, Analysis of phytoplankton yields in relation to certain physical and chemical factors of Lake Michigan: Ecology, v. 36, p. 543-552.
- Grime, E. M., 1945, Problems with iron in deep well supplies: Am. Water Works Assoc. Jour., v. 37, p. 1173-1178.
- Guelbenzu, M. D., 1951, Spectrographic analysis for trace elements: Real Acad. Farmacia Annales, v. 17, p. 237-266.
- Guseva, K. A., 1937a, The influence of manganese on the development of algae: Microbiology [U.S.S.R.], v. 6, p. 292-307.
- 1937b, Hydro- and micro-biological studies of the Ucha reservoir of the Moscow-Volga canal, part 2—The development of *Anabaena lemmermanni* Richter, *Aphanizomenon flos-aquae* and *Asterionella formosa* Hassall in the reservoir during the first summer of its existence: Microbiology [U.S.S.R.], v. 6, p. 449-464.
- 1939, The blooming of the Uchinskii reservoir: Soc. Naturalistes Moscou Bull., Sec. Biology, v. 48, no. 4, p. 30-32.
- 1940, Effect of copper on algae: Microbiology [U.S.S.R.], v. 9, p. 480-489.
- Haase, L. W., 1940, Recent views on the aggressiveness of water: Gesundheits-Ingenieur, v. 63, p. 104-107.
- Hakomori, S., 1931, Reactions of several metallic ions in the presence of organic acids: Tohoku Univ. Sci. Repts., ser., 1, v. 20, p. 736-763.
- Hale, F. E., 1949, Control of algae: Water and Sewage Works, v. 96, p. 95-96.
- Hallinan, F. J., 1943, Determination of iron in water—A modified colorimetric procedure: Indus. Eng. Chemistry, Anal. Ed., v. 15, p. 510-512.
- Hallsworth, E. G., and Costin, A. B., 1953, Pedogenesis in New South Wales, part 4—The ironstone soils: Soil Sci. Jour., v. 4, p. 24-47.
- Hance, F. E., 1934, Report on chemistry: Hawaiian Sugar Planter's Assoc. Proc., 54th Ann. Mtg., Rept. Comm.-in-charge Expt. Sta., p. 69-90.
- Handke, G., 1933, The influence of reduced iron on the aerobic bacterial flora of the intestine: Zentralbl. Bakteriologie Parasitenkunde, Abt. 1, v. 130, p. 240-247.
- Harder, E. C., 1919, Iron depositing bacteria and their geologic relations: U.S. Geol. Survey Prof. Paper 113, 89 p.
- Harley, J. L., 1940, The root system of the beech in woodland soils, with special reference to mycorrhizal infection: Jour. Ecology, v. 28, p. 107-117.
- Harmsen, G. W., 1938, Biological iron transformations in the soil: Chem. Weekblad, v. 35, p. 495-500.

- Harper, H. J., and Daniel, H. A., 1934, Chemical composition of certain aquatic plants: *Bot. Gaz.*, v. 96, p. 186-189.
- Harris, R. S., 1952, Indigenous edible plants of Latin America: *Internat. Zeitschr. Vitaminforsch.*, v. 23, p. 404-414.
- Harrison, W. H., and Subrahmanyaayyar, P. A., 1917, Note on copper sulfate as a stimulant for the rice crop: *Madras Agriculture Dept. Yearbook*, p. 55-62.
- Harvard, P. F., 1937, Minimizing iron in ground water: *Water Works Eng.*, v. 90, p. 1474-1477.
- Harvey, H. W., 1933, The rate of diatom growth: *Marine Biol. Assoc. Jour., United Kingdom*, v. 19, p. 253-276.
- 1937, The supply of iron to diatoms: *Marine Biol. Assoc. Jour., United Kingdom*, v. 22, no. 1, p. 205-219.
- 1939, Substances controlling the growth of a diatom: *Marine Biol. Assoc. Jour., United Kingdom*, v. 23, p. 499-520.
- Harward, M. E., Jackson, W. A., Lott, W. L., and Mason, D. C., 1955, Effects of aluminum, iron, and manganese upon the growth and composition of lettuce: *Am. Soc. Hort. Sci. Proc.*, v. 66, p. 261-266.
- Hasler, A., 1951, The influence of chalk on soils and upon plants: *Naturf. Gesell. Bern Mitt.*, v. 8, Sitzungsber, p. 27-29.
- Henderson, C., Robeck, G. G., and Palange, R. C., 1954, Biological aspects of low-level radioactivity in the Columbia River: *U.S. Atomic Energy Comm.*, Wash-275, p. 153-165.
- Hendrych, F., and Klimesch, K., 1935, Is there chronic iron poisoning?: *Archiv Expt. Pathologie Pharmakologie*, v. 178, p. 178-188.
- Henley, R. R., 1925, The influence of iron on the growth of the tubercle bacillus upon glycerinated beef broth: *Am. Rev. Tuberculosis*, v. 12, p. 246-259.
- Henry, C. R., 1950, Prevention of the settlement of iron: *Am. Water Works Assoc. Jour.*, v. 42, p. 887-896.
- Hermanowicz, W., and Kelus, J., 1955, Systematic technological test of water purification from iron: *Gaz., Woda i Tech. Sanitarna*, v. 29, p. 70-74.
- Hesse, P. R., 1956, Sulfate deficiency of Lake Victoria: *Nature*, v. 177, p. 389-390.
- Hester, J. B., and Shelton, F. A., 1947, Solubility of iron in submerged soils: *Science*, v. 106, p. 595.
- Hetterschij, C. W. G., 1941, The effect of temperature upon the equilibrium in soil extracts: *Chem. Weekblad*, v. 38, p. 559-561.
- Hewitt, E. J., 1950, The production of micronutrient deficiencies in plants grown in sand culture and the effects of some of these deficiencies on plant growth: *Fertilizer Soc. Proc.* 10, p. 5-12.
- 1953, The importance of trace elements in crop nutrition: *British Agr. Bull.*, v. 6, p. 222-229.
- Hewitt, L. F., 1957, Influence of hydrogen-ion concentration and oxidation-reduction conditions on bacterial behavior: *Soc. Gen. Microbiology*, 7th Symposium, p. 42-55.
- Hill, E. S., and Michaelis, L., 1933, Effect of iron on the establishment of the oxidation-reduction potential of alloxantin: *Science*, v. 78, p. 485-486.
- Hirsch, P., and Ruter, R., 1926, Reduction-oxidation potentials, part 1—Determination of minute quantities of ferrous and ferric salts.—A study of the application of reduction-oxidation potentials to analytical purposes: *Zeitschr. Anal. Chemie*, v. 68, p. 328-342.
- Hoffer, G. N., and Carr, R. H., 1920, Iron accumulation and mobility in diseased cornstalks: *Abs. in Phytopathology*, v. 10, no. 1, p. 56.

- Hofler, K., 1925, Iron content and localization of iron accumulation in the cell wall of the Desmidiaceae: Akad. Wiss. Wien., Sitzungsber, v. 135, p. 103-166.
- Holl, K., 1938, Removal of silica from raw water: Archiv Warmewirtschaft Dampfkesselwesen, v. 19, p. 323-324.
- Holland, G. J., 1956, The eradication of *Asellus aquaticus* from water supply mains: Inst. Water Engineers, Jour. v. 10, p. 221-241.
- Hollo, J., and Gorog, J., 1954, The microbial deterioration of soaps: Inst. Agr. Chem. Technology, Univ. Tech. Sci. Yearbook, Budapest, Hungary, v. 8, p. 181-183.
- Hood, J. W., 1948, Copper sulfate aids sewer maintenance and sewage treatment: Water and Sewage Works, v. 95, p. 275-277.
- Hoore, J. D., 1949, Iron compounds in the soil.—Some notes concerning their composition, role, study, and importance: Bull. Agricole Congo Belge, v. 40, p. 66-85.
- Hopkins, E. F., 1930, Iron-ion concentration in relation to growth and other biological processes: Bot. Gaz., v. 89, p. 209-240.
- 1935, Studies on the relation of iron, manganese, and copper to the growth of green plants: Cornell Univ. Agr. Expt. Sta. Bull., Ithaca, N.Y., p. 86.
- Hopkins, E. F., and Wann, F. B., 1925, The effect of the hydrogen-ion concentration on the availability of iron for *Chlorella* species: Jour. Gen. Physiology, v. 9, p. 205-210.
- 1926, Relation of hydrogen-ion concentration to growth of *Chlorella* and to the availability of iron: Bot. Gaz., v. 81, p. 353-376.
- 1927, Iron requirement for *Chlorella*: Bot. Gaz., v. 84, p. 407-427.
- Horner, C. K., and Burk, D., 1934, Magnesium, calcium, and iron requirements for growth of *Azotobacter* in free and fixed nitrogen: U.S. Dept. Agriculture Jour. Agr. Research, v. 48, p. 981-995.
- Houlihan, J. E., and Farina, P. E. L., 1953, The thiocyanate determination of iron with hydrogen peroxide: Analyst, v. 78, p. 559.
- Hseung, Y., 1941, Some chemical properties of paddy soils: China Natl. Geol. Survey, Spec. Soils Pub., 4, p. 1-22.
- Humfield, H., and Sugihara, T. F., 1952, Nutrient requirements of *Agaricus campestris* grown in submerged culture: Mycologia, v. 44, p. 605-620.
- Hutchens, J. O., 1940, The need of *Chilomonas paramecium* for iron: Jour. Cellular Comp. Physiology, v. 16, p. 265-267.
- Hutchinson, G. E., 1941, Limnological studies in Connecticut, part 4—The mechanisms of intermediary metabolism in stratified lakes: Ecol. Mon., v. 11, p. 21-60.
- 1957, A treatise on limnology in Geography, Physics, and Chemistry: New York, Wiley and Sons, v. 1, 1015 p.
- Hutchinson, G. E., Deevey, E. S., Jr., and Wollack, A., 1939, The oxidation-reduction potentials of lake waters and their ecological significance: [U.S.] Natl. Acad. Sci. Proc., v. 25, no. 2, p. 87-90.
- Hylkema, B., 1939, Portland cement as a lining for iron drinking water reservoirs (drinking water hygiene on ships at sea): Ingenieur [Utrecht], v. 54, no. 52, p. 71-75.
- Ichikawa, C., 1936, Isolation of cyanuric acid from the soil of Kagamigahara: Agr. Chem. Soc. Japan Jour., v. 12, p. 898-899.
- Ignatieff, V., 1939, Ferrous iron in soils: Nature, v. 143, p. 562-563.
- 1941, Determination and behavior of ferrous iron in soils: Soil Sci., v. 51, p. 249-263.

- Ingalls, R. A., and Shive, J. W., 1931, Relation of H-ion concentration of tissue fluids to the distribution of iron in plants: *Plant Physiology*, v. 6, no. 1, p. 103-125.
- Ingleson, H., and Thomas, G., 1946, Use of sodium hexametaphosphate in the treatment of water (containing dissolved iron) from a borehole at Scalby, Yorks: *Water and Water Eng.*, v. 49, p. 14-22.
- Inoue, S., 1932, Biological investigation of iron metabolism: *Japan Jour. Obstetrics Gynecology*, v. 15, p. 53-69.
- Isizuka, Y., 1940, The toxic action of copper ions on the growth of rice plant, part 1—The toxic action of copper ions on the growth of rice plant when half of its root is placed in solutions of copper sulfate with and without nutrients: *Jour. Sci. Soil Manure, Japan*, v. 14, p. 248-263.
- Ivanov, M. V., 1956, Using isotopes in studies of sulfate-reduction activity in Lake Belovod: *Mikrobiologiya*, v. 25, p. 305-309.
- Ivlev, V. S., 1938, The balance of substances in the "Beloe" Lake (Kosino); The balance of iron: *Limnologicheskoi Stantsii V Kosino Trudy*, 21, p. 21-59.
- Iyengar, B. A. S., 1936, Iron mobilization and plant growth in water-logged soils: *Soc. Biol. Chemists Proc., India*, v. 1, p. 32-33.
- 1937, Determination of iron in soil extracts and other biological media: *Soc. Biol. Chemists Proc., India*, v. 1, p. 11.
- Iyengar, B. A. S., and Subrahmanyam, V., 1935, Role of organic matter in plant nutrition, part 8—Influence of fermentable organic matter in the transformations of iron in swamp soil: *Indian Acad. Sci. Proc. IB*, p. 868-892.
- Izgarishev, N., and Turkovskii, A., 1929, The activation of chemical reactions by neutral salts, part 4—The effect of neutral salts on the potential of oxidation-reduction systems: *Zeitschr. phys. Chemie [Leipzig-Berlin] Abt. A*, v. 140, p. 227-234.
- Jackson, W. T., 1956, Relative importance of factors causing injury to shoots of flooded tomato plants: *Am. Jour. Botany*, v. 43, p. 637-639.
- Jacobson, L., 1945, Iron in the leaves and chloroplasts of some plants in relation to their chlorophyll content: *Plant Physiology*, v. 20, p. 223-245.
- Jacobson, L., and Oertli, J. G., 1956, Relation between iron and chlorophyll contents in chlorotic sunflower leaves: *Plant Physiology*, v. 31, p. 199-204.
- Jane, F. W., 1942, Collection and examination of fresh water algae with special reference to flagellates (microchemical methods): *Quekett Micros. Club Jour. [London, England]*, v. 1, no. 5, p. 217-219.
- Javillier, M., and Sauton, B., 1911, Is iron indispensable to the formation of Conidia of *Aspergillus niger*? : *Acad. sci. [Paris] Comptes rendus*, v. 153, no. 23, p. 1177-1180.
- Jenny, H., and Overstreet, R., 1939, Surface migration of ions and contact exchange: *Jour. Phys. Chemistry*, v. 43, p. 1185-1196.
- Joffe, J. S., 1930, Experiments on the movement of iron and aluminum in the soil: *Internat. Cong. Soil Sci.*, 2d, Leningrad, Proc., v. 2, p. 135-145.
- Joffe, J. S., and McLean, H. C., 1928, Colloidal behavior of soils and soil fertility, part 4—Anion effect on the precipitation reactions and degree of dispersion of aluminum and iron hydroxides: *Soil Sci.*, v. 26, no. 1, p. 47-59.
- Johnson, M. O., 1916, The spraying of yellow pineapple plants on manganese soils with iron sulphate solutions: *Hawaii Agr. Expt. Sta. Bull.*, v. 51, 11 p.
- Johnston, F. A., McMillan, T. G., and Evans, E. R., 1950, Perspiration as a factor influencing the requirement for calcium and iron: *Jour. Nutrition*, v. 42, p. 285-296.

- Jolivet, H., 1934, Corrosion of metals and alloys at elevated temperatures: *Metaux*, v. 9, p. 349-353.
- Juday, C., Birge, E. A., and Meloche, V. W., 1938, Mineral content of the lake waters of Northeastern Wisconsin: *Wisconsin Acad. Sci. Trans.*, v. 31, p. 223-276.
- Juliano, J. B., and Aldama, M. J., 1937, Morphology of *Oryza sativa*: *Philippine Agriculturist*, v. 26, No. 1, p. 1-134.
- Kahler, H. L., 1956, Iron retention in water supplies with water-soluble lignin: U.S. Patent No. 2,744,866.
- Kalinenko, V. O., 1940, Physiological characteristics of iron bacteria *Leptothrix ochracea*: *Acad. sci. [U.R.S.S.] Comptes rendus*, v. 29, p. 253-255.
- 1946, Role of bacteria in formation of ferromanganese concretions: *Mikrobiologiya*, v. 15, p. 364-369.
- Kapp, L. C., 1932a, Preliminary report on the effect of certain chemicals on rice production and their effect on the rice soil: *Arkansas Agr. Expt. Sta. Bull.* 277, p. 3-35.
- 1932b, Base exchange in rice soil: *Arkansas Agr. Expt. Sta. Bull.* 280, p. 23-24.
- 1934, Rice nutrition: *Arkansas Agr. Expt. Sta. Bull.* 312, p. 22-24, 33-34.
- 1938, Rice nutrition: *Arkansas Agr. Expt. Sta. Bull.* 368, p. 35-38.
- Kaserer, H., 1910, On the mineral needs of *Azotobacter*: *Deutschen Bot. Gesell. Ber.*, v. 28, no. 6, p. 208-212.
- Kato, H., 1934, Induced precipitation in chemical analysis, part 1: *Chem. Soc. Japan Jour.*, v. 55, p. 1148-1151.
- Kawamura, K., 1928, Temperature effects on the exchange acidity of lowland and swamp soils, especially the effect on the amount of acid ferrous iron: *Sci. Agr. Soc. Japan Jour.*, p. 1-20.
- Keilin, D., and Mann, T., 1944, Trace elements in relation to physiological function and enzyme systems: *Nutrition Soc. Proc.*, v. 1, p. 189-192.
- Keller, H. W., 1945, Experiences gained in furnishing water to the Army: *South-west Water Works Jour.*, v. 26, no. 10, p. 20.
- Keller, W. D., 1955, The principles of chemical weathering: Columbia, Mo., Lucas Brothers, 88 p.
- Kharasch, M. S., Conway, E. A., and Bloom, W., 1936, Some chemical factors influencing growth and pigmentation of certain microorganisms: *Jour. Bacteriology*, v. 32, p. 533-540.
- Kiene, E., 1940, The effect of trace elements on the propagation of yeast in a synthetic nutrient solution: *Vorratspflege Lebensmittelforschung*, v. 3, p. 446-456.
- Kimata, M., Kadota, H., Hata, Y., and Tajima, T., 1955, Marine sulfate-reducing bacteria, part 3—Influences of various environmental factors upon the sulfate-reducing activity of marine sulfate-reducing bacteria: *Japan Soc. Sci. Fisheries Bull.*, v. 21, p. 113-118.
- Kinne-Dietrich, E. M., 1955, The nutritive requirements of marine blue algae: *Kieler Meeresforschungen*, v. 11, p. 34-47.
- Kitano, Y., 1955, Chemical investigations of hot springs in Japan: *Chem. Soc. Japan Jour., Pure Chem. Sec.*, v. 76, p. 588-589.
- Kliman, S., 1937, The importance of ferrous iron in plants and soils: *Soil Sci. Soc. America Proc.*, v. 2, p. 385-392.
- Klut, H., 1937, The economic significance of water containing iron and manganese: *Pharmazeutische Zeitung*, v. 82, p. 881-883.

- Klyachko, V. A., 1956, Selection of a method for removal of iron compounds from the water of artesian wells: *Vodosnabzhenie Sanitarnaya Tekh.*, no. 8, p. 8-11.
- Knauss, H. J., and Porter, J. W., 1954, Absorption of inorganic ions by *Chlorella pyrenoidosa*: *Plant Physiology*, v. 29, p. 229-234.
- Knaysi, G., and Dutky, S. R., 1936, The growth of a britanol *Clostridium* in relation to the oxidation-reduction potential and oxygen content of the medium: *Jour. Bacteriology*, v. 31 (2), p. 137-149.
- Kneer, B., 1939, The purification of water containing iron, manganese, and humic acid for use in the textile industry: *Klepzig's Textilzeitschr.*, v. 42, p. 87-89.
- Koblents-Mislake, O. I., 1955, The requirement of some Black Sea algae for mineral nutritive elements: *Akad. Nauk SSSR Inst. Okeanologii Trudy*, v. 13, p. 89-93.
- Koczy, F. F., 1950, The sedimentation and geochemistry in the equatorial part of the Atlantic Ocean: *Göteborgs Kungliga Vetensk. Vitterhets-Samhälles Handl.*, v. 6, ser. B6, no. 1, 45 p.
- Koeck, H., 1956, Treatment of polluted waters by flocculation: *Annales Mines Belgique*, spec. no., p. 114-121.
- Kopal, S., 1940, Mineral and carbon dioxide-containing waters in mixtures with wines and fruit juices: *Chemicke Listy*, v. 34, p. 250-253.
- Korinek, J., 1936, The microflora of a natural copper soil: *Soc. Sci. Bohême Mém.*, v. 2, p. 1-7.
- Kosugi, T., and Umeda, K., 1931, Morphological studies of iron metabolism: *Japanese Pathology Soc. Trans.*, v. 21, p. 66-69.
- Kramer, P. J., 1957, Outer space in plants—some possible implications of the concept: *Science*, v. 125, no. 3249, p. 633-635.
- Kramli, A., Pettko, E. F., and Kiss, P., 1954, Effect of heavy metals on respiration and oxidation-reduction potentials of *Streptomyces griseus* cultures: *Acad. Sci. [Hungary] Acta Microbiol.*, v. 2, p. 39-49.
- Kraybill, H. R., 1930, Report on preparation of plant material for analysis: *Assoc. Official Agr. Chemists Jour.*, v. 13, p. 224-225.
- Kretschmer, K. F., 1952, Color measurement in wort and beer: *Brauerei Wiss. Beil.*, v. 5, no. 2, p. 15-17.
- Kriegsmann, K. F., 1938, A biological growth research on the waters of the Great Holy Lake with special reference to its iron content: *Landesmus. Provinz Westfalen, Mus. Naturkunde Abh.*, v. 9, no. 2, p. 1-106.
- Kriventsov, M. I., 1954, Concerning the expected mineralization of waters of the Stalingrad and Kubyshev reservoirs: *Gidrokhim. Materialy*, v. 22, p. 3-19.
- Kruse, H., 1949, Determination of traces of iron in water: *Gas- u. Wasserfach*, v. 90, p. 109-111.
- Kryukov, P. A., 1933, Titanometric determination of ferric iron in presence of humic matter: *Vsesoyuznyi Nauchno Issledovatel'skii Inst. Udobrenii Agrotekhnika; Agropochvovedeniya im. K. K. Gedroitsa, Trudy Leningradskogo Otdeleniya*, v. 17, p. 9-19.
- Krzemieniewski, S., and Kovats, J., 1936, The influence of iron and molybdenum on nitrogen fixation by *Azotobacter chroococcum* Beij.: *Acad. Polonaise Sci. Bull. Internat., Cl. sci. math.*, no. 8-10, p. 169-195 [1937] [in German].
- Kufferath, J., and Schmitz, D. M., 1955, Gases found in the deep waters of Lake Kivu [Belgian Congo]: *Cong. Internat. Chimie Indus. 27th, Brussels, Comptes rendus*, v. 3, *Industrie Chim. Belge*, spec. 20, p. 871-873.
- Kuhnoltz-Lordat, G., 1928, Chlorosis: *Prog. Agric. Viticole*, v. 89, p. 310-312.

- Kurzweil, H., and Exner, H., 1954, Iridescent-colored surface films of iron-containing waters: *Gesundheits-Ingenieur*, v. 75, p. 196-198.
- Lackey, J. B., 1939, Aquatic life in waters polluted by acid mine waste: U.S. Public Health Repts., v. 54, no. 18, p. 740-746.
- 1956, Stream enrichment and microbiota: U.S. Public Health Repts., v. 71, p. 708-718.
- Lackey, J. B., and Wattie, E., 1940, Pure culture studies of the sewage fungus, *Sphacrotilus natans*: U.S. Public Health Repts., v. 55, p. 975-987.
- Lahey, M. E. Gubler, C. J., Chase, M. S., Cartwright, G. E., and Wintrobe, M. M., 1952, Copper metabolism, part 3—Metabolism of iron in copper deficient swine: *Blood*, v. 7, p. 1075-1092.
- Lajoie, P. G., and De Long, W. A., 1945, The acid-oxalate extracts of podzol and podzolic soils: *Sci. Agriculture*, v. 25, p. 215-220.
- Lanyar, F., Lieb, H., and Verdino, A., 1933, The excretion of iron in human urine under physiological and pathological conditions: *Zeitschr. Physiol. Chemie*, v. 217, p. 160-166.
- Lapin, L. N., and Kill, W. E., 1931, Colorimetric estimation of iron in drinking and waste waters by means of sulfosalicylic acid: *Zeitschr. Hygiene Infektionskrankheiten*, v. 112, p. 719-723.
- Larsen, H., 1953, Microbiology and biochemistry of the photosynthetic green sulfur bacteria: *Kgl. Norske Vidensk. Seisk. Skr.*, p. 1-205.
- Lascelles, J., and Cooper, R., 1955, The formation of porphyrins and related substances by photosynthetic bacteria: *Internat. Biochimie. Cong.*, 3d, Brussels, Proc., p. 88.
- Lasseur, P., Dupaix-Lasseur, A., and Honessler, G., 1942-44, A survey of the life of microbes: *Lab. Microbiologie Fac. Pharmacie, Nancy, Travaux.*, v. 13, p. 13-54.
- Lazarev, A. M., 1939, The action of copper in marsh soils: *Khimizatsiya Sotsialisticheskogo Zemledeliya* [Chemisation Socialistic Agriculture [U.S.S.R.]] no. 7, p. 60-65.
- Leaf, W. B., 1948, Silica removal with iron shavings: *Am. Water Works Assoc. Jour.*, v. 40, p. 980-988.
- Leathen, W. W., Kinsel, N. A., and Braley, S. A., Sr., 1956, *Ferrobacillus ferrooxidans*—A chemosynthetic autotrophic bacterium: *Jour. Bacteriology*, v. 72, p. 700-704.
- Lederer, J., and Bogaert, R., 1939, Absorption of iron from chalybeate water: *Archives Med. Hydrology* [London], v. 17, p. 125-126.
- Lehmann, P., and Reuss, A., 1927, A colorimetric method for the determination of iron in water: *Zeitschr. Untersuchung Lebensmittel*, v. 54, p. 374-376.
- Leibich, H., 1941, Quantitativ-chemische untersuchungen uber das Eisen in den chloroplasten und ubrigen Zellbestandteilen von *spinacia oleracea*: *Zeitschr. Botanik*, v. 37, p. 129-157.
- Leonard, C. D., and Stewart, I., 1953, An available source of iron for plants: *Am. Soc. Horti. Sci. Proc.*, v. 62, p. 103-110.
- Levin, F. I., 1947, The migration of silica, aluminum, iron, and manganese in the profile of southern Chernozem: *Moskovskogo Univ. Vestnik* 5, p. 121-129.
- Levring, T., 1957, Growth and reproduction in marine algae in different regions: *Année Biol.*, v. 33, p. 57-65 [in English].
- Lewis, G. J. Jr., and Goldberg, E. D., 1954, Iron in marine waters: *Jour. Marine Research*, v. 13, p. 183-197.
- Lieber, M., 1953, Simplified field test for iron in water: *Water and Sewage Works*, v. 100, p. 229.

- Lieffring, P., and Buron, X., 1948, Colorimetric determination of small quantities of iron in water with dimethylglyoxime: *Chimie Anal.*, v. 30, p. 36.
- Lieske, R., 1920, Nutritional physiology of the iron bacterium: *Physiol. Abs.*, v. 5, p. 437.
- Limanowski, W., 1932, Catalytic decomposition of hydrogen peroxide by ferric ions: *Roczniki Chemii*, v. 12, p. 782-786.
- Lindquist, N., 1949, Acute iron poisoning: *Acta Paediatrica*, v. 38, p. 447-453.
- Linossier, G., 1910, The influence of iron on the formation of spores in *Aspergillus niger*: *Acad. sci. [Paris] Comptes rendus*, v. 151, no. 23, p. 1075-1076.
- Lintzel, W., 1929, Iron metabolism, part 5—The iron need of man: *Zeitschr. Biologie*, v. 89, p. 342-349.
- Lipmann, F., 1937, Photochemical reduction of ferric iron and methylene blue by pyruvic acid: *Skandinavisches Archiv Physiologie*, v. 76, p. 186-192.
- Loew, O., 1938, Intake of iron through the roots and the application of iron—Increase in yield through application of ferrous sulfate in comparison with manganous sulfate: *Angew. Botanik*, v. 20, p. 238-240.
- Lowenhaupt, B., 1955, Transport of calcium and other cations in submerged aquatic plants: U.S. Atomic Energy Comm. UCRL-3247, 40 p.
- Luhrig, H., 1927, The role of carbon dioxide in the solution of iron and manganese in water and the determination of excess carbon dioxide in water: *Gas- u. Wasserfach*, v. 70, p. 381-389.
- Lutman, B. F., 1929, *Microbiology*: New York, McGraw-Hill, p. 37-38, 53-57.
- Lutwick, L. E., Coldwell, B. B., and De Long, W. D., 1952, Leachates from decomposing leaves, part 1—Some general characteristics: *Sci. Agriculture*, v. 32, p. 99-108, 603-613.
- Lutwick, L. E., and De Long, W. A., 1954, Leachates from decomposing leaves, part 2—Interaction with soil-forming materials: *Canadian Jour. Agr. Sci.*, v. 34, p. 203-213.
- Lutz, J. F., 1936, The relation of free iron in the soil to aggregation: *Soil Sci. Soc. America Proc.*, v. 1, p. 43-45.
- Lwoff, M., 1933a, Nutrition of trypanosomes: *Annales Inst. Pasteur*, v. 51, p. 55-116.
- 1933b, Nutrition of trypanosomides and hemophilic bacteria; Baudisch's "active iron": *Annales Inst. Pasteur*, v. 51, p. 707-713.
- MacCarthy, G. R., 1925, The relationship between soluble iron and colloids in certain residual clays: *Soil Sci.*, v. 20, no. 6, p. 473-475.
- MacGillivray, J. H., Hanna, G. C., and Minges, P. A., 1942, Vitamin, protein, calcium, iron and calorie yield of vegetables per acre and per acre man-hour: *Am. Soc. Hort. Sci. Proc.*, v. 41, p. 293-297.
- Mackenzie, R. C., 1952, Investigations on cold-precipitated hydrated ferric oxide and its origin in clays—Symposium—Problems of clay and laterite genesis: New York, Am. Inst. Mining and Metall. Engineers, p. 65-75.
- MacLeod, R. A., and Onofrey, E., 1956, Nutrition and metabolism of marine bacteria, part 2—The relation of sea water to the growth of marine bacteria: *Jour. Bacteriology*, v. 71, p. 661-667.
- Magie, R. O., 1951, Soil minor-element deficiencies: *Gladiolus Mag.*, v. 15, no. 4, p. 18-21.
- Malkov, A., 1934, Role of ferrous and ferric ions in the fermentation and oxidation processes of yeasts: *Zentralbl. Bakteriologie Parasitenkunde* 2, v. 91, p. 161-163.

- Malyuga, D. P., 1945, Content of copper, nickel, cobalt, and other elements of the iron family in native waters: Acad. sci. [U.R.S.S.] Comptes rendus, v. 48, p. 113-116; Akad. Nauk SSSR Doklady, v. 48, p. 119-122.
- Mancha, R., 1936, The oxygen stratification in lakes: Archiv Hydrobiologie, v. 29, p. 695-707.
- Mandl, B., 1949, Naturally conditioned changes in the composition of the water used in brewing: Brauwelt, p. 23-25.
- Mangenot, F., 1953, The chitin of fungi: Soc. Sci. Bull., Nancy, v. 12, p. 10-18.
- Mansfield, W. W., 1953, Effect of surface films on the evaporation of water: Nature, v. 172, p. 1101.
- 1955, Influence of monolayers on the natural rate of evaporation of water: Nature, v. 175, p. 247-249.
- Maquenne, L., and Demoussy, E., 1920, Iron as toxic (to plants) and copper as antitoxic: Acad. sci. [Paris] Comptes rendus, v. 171, no. 4, p. 218-222.
- Marcellin, P., 1956, Respective role of the cuticle and of the discontinuities of the surface of apples in carbon dioxide exchange with the atmosphere: Rev. gen. botanique, v. 63, p. 193-201.
- Marchal, J. G., and Adams, R. A., 1956, Iron in the red pigment produced by *Bacillus bruntzii* and in the violet-cobalt pigment produced by *Pseudomonas cattleyae* color: Lab. Microbiologie Fac. Pharmacie, Nancy, Travaux, v. 18, p. 32-41.
- Marek, A. N., 1955, Aquatic organisms as a sanitary index of contamination of water reservoirs by radioactive substances: Gigiena Sanitariya no. 8, p. 3-9.
- Margalef, R., 1954, Determination of phytoplankton by estimation of soluble pigments and the factors affecting the relation between quantity of pigment and dry weight: Consejo Superior Inv. Cient. Pubs., Inst. Biology, v. 16, p. 71-84.
- Marsais, P., and Segal, L., 1938, The anticryptogamic action of copper: Rev. Viticulture, v. 88, p. 285-287, 305-311, 325-327.
- Marsh, R. P., 1922, Comparison of iron from some organic compounds as an essential constituent of nutrient mediums for plants: New Jersey Agr. Expt. Sta., Ann. Rept., v. 43, p. 399-402.
- Marsh, R. P., and Shive, J. W., 1925, Adjustment of iron supply to requirements of soybean in solution culture: Bot. Gaz., v. 79, no. 1, p. 1-27.
- Martell, A. E., and Calvin, M., 1952, Chemistry of metal-chelate compounds: New York, Prentice-Hall, p. 43.
- Martin, G., and Lavollay, J., 1950, The effect of copper on the permanganate titration of humic extracts—the determination of a copper index: Internat. Cong. Soil Sci., 4th, Amsterdam, Trans., v. 3, p. 83-88; v. 4, p. 78.
- Martin, J. P., 1935, Growth of sugar cane in nutrient solutions: Hawaiian Planter's Rec., v. 39, p. 79-96.
- Mason, B., 1952, Principles of geochemistry: New York, Wiley and Sons, p. 146-147.
- Masoni, G., 1914, The alkaline reaction produced in soils by acids in relation to plant nutrition, part 1—Solubility of iron compounds in the soil: Sta. Sperimentale Agraria Italy, v. 47, no. 9-10, p. 674-701.
- Mayer, A., 1892, Effect of copperas in the soil on the yield of grains: Jour. Landwirtschaft, v. 40, no. 1, p. 19-22.
- Mayer, A. M., 1952, Iron, manganese, and the reduction of nitrates by *Chlorella vulgaris*: Palestine Jour. Botany, Jerusalem, Ser., v. 5, p. 161-179.

- Mayer, A. M., and Gorham, E., 1951, The iron and manganese content of plants present in the natural vegetation of the English lake district: *Annals Botany* [London], v. 15, p. 247-263.
- Maze, P., and Perrier, A., 1904, The assimilation of certain organic substances by chlorophyll-bearing plants: *Annales Inst. Pasteur*, v. 18, no. 12, p. 721-747.
- Maze, P., Ruot and Lemoigne, 1912, Investigations on plant chlorosis induced by calcium carbonate: *Acad. sci. [Paris] Comptes rendus*, v. 155, no. 7, p. 435-437.
- McCance, R. A., and Widdowson, E. M., 1943, Iron excretion and metabolism in man: *Nature*, v. 152, p. 326-327.
- McDonald, J. A., 1935, Phosphate fixation in soils in relation to iron availability and its possible connection with the gypsum-phosphate problem in cacao soils: Imperial College of Tropical Agriculture, Trinidad, 4th Ann. Rept. Cacao Research, p. 86-87.
- McGeorge, W. T., 1939, Some problems connected with fertilization of alkali soils: *California Citrograph*, v. 24, no. 11, p. 389, 424.
- 1951, Lime-induced chlorosis in western crops: Better crops with plant food, v. 35, no. 5, p. 17-20, 43-46.
- McGeorge, W. T., and Breazeale, B. I., 1956, Application of the Neubauer technique and applied potential to the study of immobilization of iron in plants: *Soil Sci.*, v. 82, p. 329-336.
- McIntire, F. C., Riker, A. J., and Peterson, W. H., 1941, The role of certain vitamins and metallic elements in the nutrition of the crown-gall organism: *Jour. Bacteriology*, v. 42, p. 1-13.
- McMurtrey, J. E., Jr., 1933, Distinctive effects of the deficiency of certain essential elements on the growth of tobacco plants in solution cultures: U.S. Dept. Agriculture Tech. Bull. 340, 43 p.
- 1952, Deficiencies of secondary and micronutrient elements in plants: Better crops with plant food, v. 36, no. 9, p. 19-26, 42-48.
- Meehan, W. J., and Baas-Becking, L., 1927, Iron organisms: *Science*, v. 66, p. 42.
- Melnichenko, Z. A., 1954, Oxidation-reduction processes in the leaves and roots of plants: Gor'kovskogo Gosudarstvennogo Univ. Uchenye Zapiski, 25, p. 77-85.
- Messikommer, E., 1956, Algae of the eastern Bernese oberland: *Naturf. Gesell. Bern Mitt.*, v. 13, p. 81-152 [in German].
- Metz, O., 1930, Effect of iron, zinc, and copper on the growth and pigmentation of molds: *Archiv Mikrobiologie*, v. 1, p. 197-250.
- Metzger, W. H., 1930, Replaceable bases of irrigated soil: *Soil Sci.*, v. 29, p. 251-260.
- Meyer, R., and Bunger, J., 1952, The determination of nitrogen compounds and iron in smallest amounts of drinking water with Pulfrich's photometer: *Gesundheits-Ingenieur*, v. 73, p. 367-372.
- Midgley, A. R., 1940, Phosphate fixation in soils—a critical review: *Soil Sci. Soc. America Proc.*, v. 5, p. 24-30.
- Milad, Y., 1939, Physiological studies in lime-induced chlorosis: Ministry Agriculture Egypt Tech. Sci. Service Bull. 211, 56 p.
- Mills, E. V., 1932, Ultimate analysis of sewage: *Soc. Chem. Industry Jour. [London]*, v. 51, p. 205-209 T.
- Milne, L. J. and Milne, M. J., 1952, The biotic world and man: New York, Prentice-Hall, 588 p.

- Minami, K., Hyodo, S., and Yamane, I., 1955, Growth and respiration of tubercle bacteria: *Fukushima Jour. Med. Sci.*, v. 2, p. 99-120.
- Mitchell, C. A., 1944, Standardization of the tannin used for the comparison in Nierenstein's micromethod of estimating tannin: *Analyst*, v. 69, p. 92.
- Mohler, H., 1951, Iron and manganese as disturbing factors in ground water: *Schweizer. Ver. Gas- u. Wasserfachmännern Monats-Bull.*, v. 31, p. 293-299, 322-331.
- Monnier, A., and Kuczraski, L., 1917, The use of iron in agriculture: *Archives Sci. Phys. Nat.*, v. 43, no. 1, p. 66-68.
- Moore, B., 1914, The presence of inorganic iron compounds in the chloroplasts of the green cells of plants considered in relationship to natural photosynthesis and the origin of life: *Royal Soc. [London] Proc., Ser. B.* 87, no. B 598, p. 556-570.
- Moore, D. P., Harward, M. E., Mason, D. D., Hader, R. J., Lott, W. L., and Jackson, W. A., 1957, Some of the relationships among copper, iron, and molybdenum in the growth and nutrition of lettuce, part 2—Response surfaces of growth and accumulations of copper and iron: *Soil Sci. Soc. America Proc.*, v. 21, p. 65-74.
- Moore, W. G., 1950, Limnological studies of Louisiana lakes, part 1—Lake Providence: *Ecology*, v. 31, p. 86-99.
- 1952, Limnological studies of Louisiana lakes, part 2—Lake Chicot: *Louisiana Acad. Sci. Proc.*, v. 15, p. 37-49.
- Morello, M. A., 1940, Microchemical detection and distribution of iron in plants: *Mikrochemie vereinigt Mikrochimica Acta*, v. 28, p. 245-253.
- Morgan, M. F., 1941, Chemical soil diagnosis by the universal soil testing system: *Connecticut Agr. Expt. Sta. [New Haven] Bull.* 450, p. 579-628.
- Morison, C. G. T., and Doynne, H. C., 1914, Ferrous iron in soils: *Jour. Agr. Sci. [England]*, v. 6, no. 1, p. 97-101.
- Morris, R. L., 1952, Determination of iron in water in the presence of heavy metals: *Anal. Chemistry*, v. 24, p. 1376-1378.
- Mortimer, C. H., 1941, The exchange of dissolved substances between mud and water in lakes: *Jour. Ecology*, v. 29, p. 280-329.
- 1942, The exchange of dissolved substances between mud and water in lakes: *Jour. Ecology*, v. 30, p. 147-201.
- Moss, M. I., and Mellon, M. G., 1942, Colorimetric determinations of iron with 2,2'-bipyridyl and with 2,2',2'' terpyridyl: *Indus. Eng. Chemistry, Anal. Ed.*, v. 14, p. 862-865.
- Muenschel, W. C., 1944, *Aquatic plants of the United States*: Ithaca, N.Y., Comstock Publishing Co., 374 p.
- Mukherjee, J. N., and Chatterjee, B., 1945, Liberation of H^+ , Al^{+++} , and Fe^{+++} ions from hydrogen clays by neutral salts: *Nature*, v. 155, p. 268-269.
- Mukherjee, J. N., Chatterjee, B., and Banerjee, B. M., 1947, Liberation of hydrogen, aluminum, and ferric ions from hydrogen clays by neutral salts: *Jour. Colloid Sci.*, v. 2, p. 247-256.
- Mukherjee, J. N., Chatterjee, B., and Ray, A., 1946, Liberation of H^+ , Al^{+++} , and Fe^{+++} ions from pure clay minerals on repeated soil treatment and desaturation: *Jour. Colloid Sci.*, v. 3, p. 437-446.
- Muller, H., 1933, Limnological field methods: *Internat. Rev. gesamten Hydrobiologie Hydrographie*, v. 28, p. 351.
- Muller, J., 1950, Enormous variations in the iron content of a deep well water and an explanation: *Gas- u. Wasserfach*, v. 90, p. 623-624.

- Mulwani, B. T., and Pollard, A. G., 1939, The practical significance of soil solution studies: *Indian Jour. Agr. Sci.*, v. 9, p. 473-502.
- Mundkur, B. B., 1928, Importance of iron for the growth of fungi (*Fusarium vasinfectum* Atkinson): *Indian Sci. Cong.*, 15th Proc. p. 230.
- Murray, J., 1908, The distributions of organisms in the hydrosphere as affected by varying chemical and physical conditions: *Internat. Rev. gesamten Hydrobiologie Hydrographie*, v. 1, p. 10-17.
- Murray, R. C., 1956, Recent sediments of three Wisconsin lakes: *Geol. Soc. America Bull.*, v. 67, p. 883-910.
- Muth and Voigt, 1929, The phosphoric acid and iron content of Rhengau musts and wines: *Landwirtschaftliche Jahrbücher*, v. 68, supp. 1 p. 422-426.
- Myers, J., Phillips, J. N., and Graham, J. R., 1951, On the mass culture of algae: *Plant Physiology*, v. 26, p. 539-548.
- Nakabayashi, T., 1954, Tannins of *Spirogyra arcta*, part 4—Estimation of tannin in *Spirogyra arcta*: *Agr. Chem. Soc. Japan, Jour.*, v. 28, p. 958-961.
- Narasinhmurthy, G. 1940, Carbon transformation during the decomposition of cane molasses under swamp soil conditions: *Indian Jour. Agr. Sci.*, v. 10, p. 768-772.
- Narasinhmurthy, G., and Subrahmanyam, V., 1935, Role of organic matter in plant nutrition, part 7—Economy of carbon during decomposition of cane molasses in the swamp soil: *Indian Acad., Sci. Proc. 1B*, p. 823-836.
- Naumann, E., 1928a, The term iron-organism: *Deutsche Bot. Gesell. Ber.*, v. 46, p. 135-140.
- 1928b, Iron-organisms and the formation of lake ore deposits—certain fundamental aspects of this question: *Deutsche Bot. Gesell. Ber.*, v. 46, p. 141-147.
- Navellier, P., 1950, Iron in foods: *Annales Falsifications Fraudes*, v. 43, p. 361-363.
- Nave, P., 1947, Particular properties of the hexametaphosphate ion: *Technique l'eau [Brussels]*, p. 11-12.
- Needham, J. G., and Needham, P. R., 1935, A guide to the study of fresh-water biology: Ithaca, N.Y., Comstock Publishing Co., Inc., 88 p.
- Nehira, T., 1955, Sporulation of *Aspergillus oryzae*, part 4—Comparison of various media: *Jour. Fermentation Technology*, v. 33, p. 313-317.
- Neillands, J. B., 1957, Some aspects of microbial iron metabolism: *Bacteriological Rev.*, v. 21, p. 101-111.
- New Jersey State Agricultural Experiment Station, 1928, Plant physiology: New Jersey State Agr. Expt. Sta. Rept., p. 58-61.
- New Process Rayon Inc., 1938, Removing iron from copper: New Process Rayon Inc., French Patent No. 827,233.
- Newell, B. S., 1957, The nocturnal reduction of dissolved oxygen in dams: *East African Agr. Jour.*, v. 23, p. 127-129.
- Newell, J. M., and McCollum, E. V., 1931, Spectrographic analysis of marine products: U.S. Dept. Commerce, Bur. Fisheries, Investigational Rept. v. 5, p. 1-9.
- Nicholas, D. J. D., 1949, The manganese and iron contents of crop plants as determined by chemical methods: *Jour. Hort. Sci.*, v. 25, p. 60-77.
- Nicholas, D. J. D., and Commission, K., 1957, Effects of molybdenum, copper, and iron on some enzymes in *Neurospora crassa*: *Jour. Gen. Microbiology*, v. 17, p. 699-707.
- Nierenstein, M., 1945, Tannin content of the *Potamogeton* species and its effect on the color of iron-containing inland waters: *Analyst*, v. 70, p. 427.

- Nietsch, B., 1956, Mechanism of repeated iron pickup by waters: Gas, Wasser, Wärme, v. 10, p. 224-225.
- Nikitin, A. A., 1950, Value of rare elements: Commercial Fertilizer, v. 80, no. 4, p. 20-29.
- Nishida, H., 1954, Rapid colorimetric determination of traces of iron in water: Japan Analyst, v. 3, p. 250-251.
- Nitschmann, H., 1938, Reaction direction by nuclei—the oxidation of iron in ferrous sulfate solution: Helvetica Chim. Acta, v. 21, p. 1609-1618.
- Nordell, E., 1953, Iron and manganese removal: Water and Sewage Works, v. 100, no. 5, p. 181-185.
- Notarov, V. D., 1956, The formation conditions of sodium chloride-containing waters of the Krivbass (Krivorozhsk iron ore deposits): Akad. Nauk Ukrainskoi RSR, Geol. Zhur., v. 16, no. 3, p. 38-42.
- Novelli, A., 1953, Factors influencing production of pigment by *Serratia marcescens*—Role of iron and certain pyrrole groups: Ist. Sieroterapico Milanese Boll., v. 32, p. 323-326.
- Novozhilova, M. I., 1955, Kinetics of bacterial cell counts and biomass in water layers in the Rybin reservoir: Mikrobiologiya, v. 24, p. 710-717.
- Oesting, R. B., 1934, A modified Van Slyke method for the determination of oxygen and carbon dioxide in water: Physiol. Zool., v. 7, p. 542-549.
- Ohira, K., 1955, Absorption and translocation of phosphorus and iron in rice plants: Tohoku Jour. Agr. Research, v. 5, p. 345-353.
- Ohle, W., 1937, Kolloidgele als Nahrstoffregulatoren der Gewässer: Naturwissenschaften, v. 25, p. 471-474.
- Okura, T., and Goto, K., 1955, Oxidation of ferrous ion in natural water: Chem. Soc. Japan Jour., Indus. Chem. Sec., v. 58, p. 239-241.
- Okura, T., and Kayama, I., 1952, Changes (of composition) in natural water: Kagaku [Tokyo,] v. 22, p. 262-263.
- Olivier, H. R., and Le Peintre, M., 1955, Solubilization by bio-chemical means of arsenic bound to iron in the soil: Annales Inst. Pasteur, v. 88, p. 668-671.
- Olsen, C., 1930, The influence of humus substances on the growth of green plants in water culture: Lab. Carlsberg, Comptes rendus Travaux Sér. chim., v. 18, p. 1-16.
- 1935, Iron absorption and chlorosis in green plants: Lab. Carlsberg, Comptes rendus Travaux, Sér. chim., v. 21, p. 15-52.
- 1938, Experiments with different quantities of iron salts given to maize in water culture: Lab. Carlsberg, Comptes rendus Travaux Sér. chim., v. 21, no. 23, p. 301-313.
- Olsen, S., 1950, Aquatic plants and hydrospheric factors: Svensk Bot. Tidskr., v. 44, no. 1, p. 1-31.
- Olson, R. A., Brust, H. F., and Tressler, W. L., 1941, Studies of the effects of industrial pollution in the lower Patapsco River area, part 1—The Curtis Bay Region: Chesapeake Biology Lab. Pub. 43, 40 p.
- Olson, R. V., 1947, Iron solubility in soils as affected by pH and free iron oxide content: Soil Sci. Soc. America Proc., v. 12, p. 153-157 [1948].
- O'Meara, R. A. Q., 1937, Some factors which influence reduction in broth and their bearing on the growth of *Clostridium tetani*: Jour. Pathology Bacteriology, v. 45, p. 541-553.
- Ono, N., 1900, The influence of chemical agents on the growth of algae and fungi: Imp. Univ. Tokyo Coll. Sci. Jour., v. 13, p. 141-186.

- Onodera, I., 1928, Studies on the decomposition of genge (*Astragalus sinicus*) in the soil and its relation to the growth of paddy rice: Pan-Pacific Sci. Cong., 3d, 1926, Proc. 2, p. 1973-1988.
- Ornstein, G., 1935, Preparation of water for the textile industry, removal of algae and sterilization: Melliland Textilber., v. 16, p. 760.
- Osugi, V. S., and Aoki, M., 1936, Abnormally acid soil, part 4—Existence of organic acids in the soil: Agr. Chem. Soc. Japan Jour., v. 12, p. 995-1003.
- 1938a, Abnormal acid soil: Agr. Chem. Soc. Japan Jour., v. 14, p. 251-262.
- 1938b, On the acid soil of peculiar behavior: Kyoto Coll. Agriculture Mem., v. 45, Chem. Ser. 23, 68 p.
- Osugi, V. S., Aoki, S., and Morita, S., 1934, Peculiar acidic soils: Agr. Chem. Soc. Japan Jour., v. 10, p. 443-450.
- 1935, Abnormal acid soil, part 2: Agr. Chem. Soc. Japan Jour., v. 11, p. 34-49.
- Osugi, V. S., Nishigaki, N., and Yoshimi, M., 1935, Abnormal acidic soils, part 3: Agr. Chem. Soc. Japan Jour., v. 11, p. 659-673.
- Oudin, M., 1938, Iron and pedogenic processes: Congrès de chimie industrielle, Compte rendu analytique Cong. Chimie Indus., 18th, Nancy, Comptes rendus, p. 134-137.
- Ouellette, G. J., 1951, Iron-manganese interrelationships in plant nutrition: Sci. Agriculture, v. 31, p. 277-285.
- Oy, E., 1940, Content of iron, copper, manganese, and boron in seaweeds and kelp: Tidsskr. Kjem. Bergvesen, v. 20, p. 114-117.
- Palit, C. C., Basu, S. N. and Varma, V. L., 1944, Epidemic dropsy and mustard oil: Univ. Allahabad Studies, Chemistry Sec., p. 1-10.
- Parker, C. A., 1955, Anaerobiosis with iron wool: Australian Jour. Expt. Biology Med. Sci., v. 33, p. 33-38.
- Parks, T. B., and Rose, E. R., 1933, Copper, iron, and manganese content of fish: Jour. Nutrition, v. 6, p. 95-98.
- Parsche, F., 1940, Lime chlorosis of lupines, part 2: Bodenkunde u. Pflanzenernährung, v. 19, p. 55-80.
- Patrick, W. A., 1953, Deoxidizing water: U.S. Patent No. 2,663,688.
- Pearsall, W. H., 1950, The investigation of wet soils and its agricultural implications: Empire Jour. Expt. Agriculture, v. 18, p. 289-298.
- 1952, The pH of natural soils and its ecological significance: Jour. Soil Sci., v. 3, no. 1, p. 39-51.
- Perosa, L., Dell'Aquila, M. D., and Bini, L., 1951, Elimination of iron with sweat: Soc. Italiana Biologia Sperimentale Boll., v. 27, p. 919-921.
- Peterson, E. A., and Katznelson, H., 1954, The nutrition of *Helminthosporium sativum* and certain related species: Canadian Jour. Microbiology, v. 1, p. 190-197.
- Peterson, W. H., and Elvehjem, C. A., 1928, The iron content of plant and animal foods: Jour. Biol. Chemistry, v. 78, no. 1, p. 215-223.
- Piery, M., Enselle, J., and Hery, M., 1938, The mode of action of natural water containing iron: Acad. Medicine Bull., v. 120, p. 774-778.
- Pika, L., 1956, The polarity of copper and the iron oxide Fe_2O_3 in alkaline solutions: Mitt. Vereinigung Grosskesselbesitzer, p. 433-434.
- Pillai, S. C., 1938, A biochemical investigation of the tuberculation of water pipes: Natl. Inst. Sci. India Proc., v. 4, p. 295-318.
- Pirrie, R., 1950, Iron metabolism—A review of recent literature: Glasgow Med. Jour., v. 31, p. 397-416.

- Pobeguín, T., 1954, The carbonates of calcium—Precipitation of calcium by plants compared with that in the animal world: *Annales Sci. Nat., Botanique Biologique Vegetale*, v. 15, p. 29–109.
- Polster, M., Rohan, P., and Simikova, V., 1953, Influence of iron on growth of mycobacteria: *Ceskoslovenska Hygiena Epidemiologie Mikrobiologie Imunologie*, v. 2, p. 424–431.
- Pomeroy, R., 1942, The determination of iron in sewage: *Sewage Works Jour.*, v. 14, p. 651–652.
- Pond, R. H., 1903, The biological relation of aquatic plants to the substratum: U.S. Commissioners Fish and Fisheries Rept., v. 117, p. 483–526.
- Ponnamperuma, F. N., Bradfield, R., and Peech, M., 1955, Physiological disease of rice attributable to iron toxicity: *Nature*, v. 175, p. 265.
- Porter, L. K., and Thorne, D. W., 1955, Interrelation of carbon dioxide and bicarbonate ions in causing plant chlorosis: *Soil Sci.*, v. 79, p. 373–382.
- Posnjak, E., 1926, Acceleration of rate of oxidation of ferrous iron in presence of copper, and its application to the “heap leaching” process: *Am. Inst. Mining Metall. Eng.*, no. 1615D, 10 p.
- Possingham, J. V., and Brown, R., 1957, Intracellular incorporation of iron 59 into the root cells of *Pisum*: *Nature*, v. 180, p. 653–654.
- Prescott, G. W., 1956, A guide to the literature on ecology and life histories of the algae: *The Bot. Rev.*, v. 22, no. 3, p. 167–240.
- Pringsheim, E. G., 1934, Investigations on Uspenski's iron hypothesis of the distribution of algae: *Planta*, v. 22, p. 268–312.
- 1946, Iron flagellates: *Royal Soc. [London] Trans.*, v. B232, p. 311–342.
- 1949, Iron bacteria: *Biol. Rev.*, v. 24, p. 200–245.
- Prokhorova, Z. A. 1957, Dynamics of nutritive conditions and of oxidation reduction processes in the Moscow River bottomlands: *Pochvovedenie* 1, p. 52–61.
- Pronin, M. E., 1933, The behavior of ferrous iron in normal soils: *Severo Kavkazskii Zernovoi Inst. (North Caucasian Grain Inst.)*, Colln. Sci. Paper 1, p. 114–120.
- Prozorovskaya, A. A., 1936, The influence of humic acid and its derivatives on the intake of nitrogen, phosphorus, potassium and iron by plants: *Sci. Inst. Fertilizers Insectofungicides [U.S.S.R.] Trans* 127, p. 143–161.
- Puri, A. N., and Sarup, A., 1938, Oxidation-reduction potentials in soils: *Soil Sci.*, v. 46, p. 323–329.
- Puustjarvi, V., 1952, The precipitation of iron in peat soils: *Acta Agr. Fennica*, v. 78, p. 3–72.
- Puustjarvi, V., and Juusela, T., 1952, Rust precipitates present in drainage pipes and the means of preventing their formation: *Acta Agr. Scandinavica*, v. 2, p. 131–152.
- Quartaroli, A., 1932, The iron-copper couple in chemistry and biology: *Annali Chimica Appl.*, v. 22, p. 517–543.
- Rabotnova, I. L., Toropova, E. G., and Rabaeva, M. Y., 1955, Requirements of anaerobic bacteria in the oxidation-reduction conditions of mediums: *Mikrobiologiya*, v. 24, p. 525–531.
- Rackow, B., 1957, Is photosynthesis in plants a reaction in light with an opposing dark reaction? A trial of the cyclic process with ferrous carbonate: *Planta*, v. 49, p. 494–497.
- Rakestraw, N. W., Mahneke, H. E., and Beach, E. F., 1936, Determination of iron in sea water: *Indus. Eng. Chemistry, Anal. Ed.*, v. 8, p. 136–138.
- Ramaut, J., 1955, Origin of acidity of the peat bogs of Baraque-Michel: *Acad. royale Belgique Bull.*, cl. sci., v. 41, p. 1037–1052.

- Randoin, L., and Fournier, P., 1947, Studies of the phosphorus, calcium, and iron content of the leaves of *Petroselinum sativum*, a source exceptionally rich in calcium and iron: Soc. Sci. Hygiene Alimentaire Bull., v. 35, p. 184-193.
- Randoin, L., and Le Gallic, P., 1942, Existence of correlations between the protein, iron, and vitamin contents of plant tissues: Soc. Chimie Biol. Bull., v. 22, p. 593-602.
- Rankama, K., and Sahama, T. G., 1952, Geochemistry: Chicago, Ill., Univ. Chicago Press, p. 662-676.
- Read, B. E., and Gow, G. K., 1927, Iodine, arsenic, iron, calcium, and sulfur content of Chinese medicinal algae: Chinese Jour. Physiology, v. 1, p. 99-108.
- Reed, H. S., and Haas, A. R. C., 1924, Iron supply in a nutrient solution: Bot. Gaz., v. 77, no. 3, p. 290-299.
- Reilhes, R., 1944, Cytological study of penetration of mineral substances into roots: Soc. biologie Paris Comptes rendus, v. 138, p. 316-317.
- Reisner, G. S., and Thompson, J. F., 1956, Large scale laboratory culture of *Chlorella* under conditions of micronutrient element deficiency: Plant Physiology, v. 31, p. 181-185.
- Remezov, N. P., 1929, Oxidation and reduction processes in podzol soil: Zeitschr. Pflanzenernahrung Dungung 15A, p. 34-44.
- 1930, The dynamics of oxidation-reduction potentials in podzol soils: Sci. Inst. Fertilizers [Moscow] Trans. 77, p. 45-63.
- Remington, R. E., and Shiver, H. E., 1930, Iron, copper, and manganese content of some common vegetable foods: Assoc. Official Agr. Chemists Jour., v. 13, no. 1, p. 129-132.
- Richter, O., 1926, The iron requirement of rice plants: Akad. Wiss. Wien Sitzungsber., Math-naturw. Kl., v. 135, I, nos. 5-6, p. 203-242.
- Rideal, E. K., 1925, On the influence of thin surface films on the evaporation of water: Jour. Phys. Chemistry, v. 29, p. 1585-1588.
- Riley, G. A., 1939, Limnological studies in Connecticut: Ecol. Mon., v. 9, p. 53-94.
- 1940, Limnological studies in Connecticut, part 3—The plankton of Linsley, Pond: Ecol. Mon., v. 10, p. 279-306.
- Rippel, A., 1923, The iron chlorosis in green plants caused by manganese: Biochem. Zeitschr., v. 140, p. 315-323.
- 1941, Metal catalysis and the production of humus-like substances: Forschungsdienst, v. 17, p. 54-58.
- Ritter, W., and Christen, M., 1935, Fishiness in butter, part 2—The metal content of cream and butter: Schweizer. Milchzeitung, v. 61, p. 23-24.
- Roberg, M., 1932, The physiology of assimilation in green algae, part 2—The effect of iron, zinc, and copper salts: Jahrb. Wiss. Bot., v. 76, p. 311-332.
- Robinson, W. O., 1930, Some chemical phases of submerged soil conditions: Soil Sci., v. 30, p. 197-217.
- 1951, The minor or trace elements in soils, plants, and animals: Beltsville, Md., U.S. Bur. Plant Industry Soils Agr. Eng., Agr. Research Service, 9 p.
- Rodriguez, C., and Abadia, A., 1956, The rooting of quince trees as affected by induced iron deficiency: Anales Edafologia Fisiologia Vegetal [Madrid], v. 15, p. 161-166 [English summary].
- Roelofs, E. W., 1940, Available plant nutrients in lake soils: Michigan State Coll. Quart. Bull. 22, no. 4, p. 247-254.
- 1944, Water soils in relation to lake productivity: Michigan Agr. Expt. Sta. Tech. Bull. 190, 31 p.

- Rozenberg, L. A., and Mefedova, N. A., 1956, Relations of bacteria to chemical processes and animal organisms in sea muds of the Northwest Pacific: *Mikrobiologiya*, v. 25, p. 61-65.
- Rudge, E. A., 1936, The decomposition of timber under industrial conditions, part 8—The effect of calcium and iron: *Soc. Chem. Indus. Jour.*, v. 55, p. 179-180.
- Rudolfs, W., 1922, Oxidation of iron pyrites by sulfur oxidizing organisms and their use for making mineral phosphates available: *Soil Sci.*, v. 14, no. 2, p. 135-147.
- Rummenl, G., 1955, Model studies of the physiological oxidation-reduction potential of plants and the chemism of its regulation: *Friedrich-Schiller-Univ. Jena Wiss. Zeitschr., Math. naturw. Reihe*, p. 611-632.
- Runehjelm, D., 1930, Micro-determination of iron in leaves with deficient chlorophyll: *Biochem. Zeitschr.*, v. 224, p. 481-486.
- Saeger, A. C., 1937, The concentration of copper in nutrient solutions for *Spirodelia polyrhiza*: *Am. Jour. Botany*, v. 24, p. 640-643.
- Saint, S. J., 1935, Report of chemical section (sulfate, copper, and iron in molasses): *Agr. Jour. [Barbados]*, v. 4, p. 66-73.
- Samish, R. M., 1954, Lime-induced chlorosis in grape vines: *Cong. Internat. Bot.*, 8th, Paris, p. 156-165.
- Sandstrom, A. E., and Von Gegerfelt, M., 1945, Traces of iron in the glacier ice: *Arkiv Matematik, Astronomi, Fysik, [Stockholm]*, v. B32, no. 6, 4 p. [in English].
- Saraswathi-Devi, L., 1955, Heavy-metal requirements of fungi: *Indian Acad. Sci. Proc.*, 41B, p. 145-150.
- Sarishvili, I. F., and Bagaturiya, E. A., 1938, Indications of the absorption of phosphoric acid (by soils): *Sovetskie Subtropiki*, no. 2, p. 70-72.
- Saruhashi, K., 1955, Metabolism in natural waters, part 2—Equilibrium, concentration ratio of carbonic acid substances dissolved in natural water: *Meteorol. and Geophys. [Japan] Papers*, v. 6, p. 38-55.
- Saunders, D. J., 1938, Removal of iron and hardness from water: *New England Water Works Assoc. Jour.*, v. 52, p. 443-451.
- Saunders, G. W., 1957, Interrelations of dissolved organic matter and phytoplankton: *Bot. Rev.*, v. 23, p. 389-409.
- Sauton, B., 1911, Influence of iron on the growth of certain molds: *Annales Inst. Pasteur*, v. 25, no. 12, p. 922-928.
- Sayre, J. D., 1930, Accumulated iron in the nodes of corn plants: *Plant Physiology*, v. 5, p. 393-398.
- Schachtschabel, P., 1942, Investigations of sorption in soils: *Forschungsdienst*, v. 16, p. 87-90.
- Scharrer, K., and Schropp, W., 1933, Potassium-iron antagonism in plants: *Zeitschr. Pflanzenernährung Düngung Bodenkunde*, v. 28A, 1933, p. 158-172.
- Schatz, V., Schatz, A., Trelawny, G. S., and Barth, K., 1956, Significance of lichens as pedogenic (soil-forming) agents: *Pennsylvania Acad. Sci. Proc.*, v. 30, p. 62-69.
- Schenck, R., Dingmann, T., Kirscht, P. H., and Kortengraber, A., 1932, Equilibrium investigations of the reduction, oxidation, and carbonization process of iron, parts 10 and 11—The system—Iron-oxygen, abnormal oxidation; activation of iron without additions: *Zeitschr. Anorg. Allg. Chemie*, v. 206, p. 73-96.
- Scheringa, K., 1931, Colorimetric iron determinations in drinking water and in chalybeate preparations: *Pharmaceutische Weekblad*, v. 68, p. 735-738.

- Schilling, K., 1936a, Sampling and analysis of water: Gas- u. Wasserfach, v. 79, p. 229-231.
- 1936b, Aggressive waters and their treatment: Gas- u. Wasserfach, v. 79, p. 542-545.
- Schimper, A. F. W., 1903, Plant geography upon a physiological basis: Oxford, England, Clarendon Press, 839 p.
- Schmidt-Nielsen, S., 1938, The toxicity to fish of sewage containing copper and iron: Kgl. Norske Vidensk. Selsk. Förh., v. 11, p. 233-236.
- Schnitzer, M., and De Long, W. A., 1954a, Note on the reaction of 2,2' dipyridyl with iron in the presence of organic matter: Canadian Jour. Agr. Sci., v. 34, p. 324-325.
- 1954b, Note on relative capacities of solutions obtained from Forest vegetation for mobilization of iron: Canadian Jour. Agr. Sci., v. 34, p. 542-543.
- 1955, Investigations on the mobilization and transport of iron forested soils, part 2—The nature of the reaction of leaf extracts and leachates with iron: Soil Sci. Soc. America Proc., v. 19, no. 3, p. 363-368.
- Schoeller, M., 1955, The solubility of iron in subterranean waters: Annales Inst. Hydrologie Climatologie, v. 26, p. 1-37.
- Schoeller, W. R., and Webb, H. W., 1936, Investigations into the analytical chemistry of tantalum, columbium, and their mineral associates, part 32—Observations on phosphorus, vanadium, and a tannin precipitation series: Analyst, v. 61, p. 585-590.
- Scholz, W., 1932, Microcolorimetric determination of iron in small amounts of plant ash: Zeitschr. Pflanzenernährung Düngung Bodenkunde, v. 26A, p. 212-216.
- Schroder, H., 1939, Die Algenflora der Mulde Ein Beitrag zur Biologie saprober Flüsse: Pflanzenforschung, v. 21, 88 p.
- Schwaibold, J., and Fischler, F., 1931, The reciprocal action between metal utensils and foodstuffs, part 2—A study of the biological value of metals: Biochem. Zeitschr., v. 232, p. 240-253.
- Schwartz, M., 1956, The reduction of ferricyanide by living *Chlorella* cells: Archives Biochemistry Biophysics, v. 63, p. 201-211.
- Scott, J. P., and Brandley, C. A., 1933, The use of reduced iron for the cultivation of anaerobic organisms: Jour. Bacteriology, v. 26, p. 1-7.
- Scouller, W. D., 1932, Effect of iron on sewage purification: Surveyor, v. 82, p. 476.
- Seeberg, J., 1937, The ferrous-ion concentration of ground water and its variation: Univ. Latviensis, Acta, Lauksaimniecibas Fakultat Ser., v. 3, no. 4-7, p. 325-419. [in Lettish.]
- Segeberg, H., 1937, The acidity of marshy soils—in particular that due to iron disulfide and its determination: Bodenkunde u. Pflanzenernährung, v. 4, p. 50-64.
- Serdobolskii, I. P., 1950a, Oxidation-reduction and alkaline-acid conditions for glei formation: Akad. Nauk SSSR, Trudy Pochvennogo Inst. V. V. Dokuchaeva, v. 31, p. 73-81.
- 1950b, The influence of soil conditions on the transformation of manganese compounds in the soil: Akad. Nauk SSSR, Trudy Pochvennogo Inst. V. V. Dokuchaeva, v. 33, p. 192-216.
- Serdobolskii, I. P., and Shavrygin, P. I., 1950, Oxidation-reduction conditions of solanchak meadow soils of the Fergan Valley: Akad. Nauk SSSR, Trudy Pochvennogo Inst. V. V. Dokuchaeva, v. 31, p. 82-90.

- Setter, L. R., 1930, Effect of iron on the anaerobic decomposition of sewage sludge: *Sewage Works Jour.*, v. 2, p. 504-520.
- Sharpe, L. M., 1948, Radioactive iron as a biological tracer: *Brookhaven conf. Rept. BNL-C-4*, p. 126-136.
- Shcherbina, V., 1939, Oxidation-reduction potentials as applied to the study of the paragenesis of minerals, part 6: *Acad. sci. [U.R.S.S.]. Comptes rendus*, v. 22, p. 503-506 [in English].
- Sheets, O., Leonard, O. A., and Gieger, M., 1941, Distribution of minerals and vitamins in different parts of leafy vegetables: *Food Research*, v. 6, p. 553-569.
- Sherman H. C., 1907, Iron in food and its function in nutrition: *U.S. Dept. Agriculture, Office Expt. Sta., Bull. 185*, 80 p.
- Shidlovskaya-Ovchinnikova, Yu S., 1953, Stabilizing ferrous iron in natural waters for the purpose of determining separately the ferrous and ferric iron: *Gidrokhimi. Materialy*, v. 20, p. 79-91.
- Shilova, E. I., and Kreier, K. G., 1957, Carbonic acid of the soil solution and its role in soil formation: *Pochvovedenie*, 7, p. 65-72.
- Shkol'nik, M. Y., and Makarova, N. A., 1950, Antagonism of iron and copper: *Doklady Akad. Nauk SSSR*, v. 70, p. 121-124.
- Shkolnik, M. V., and Steklova, M. M., 1954, The nature of some similarity of action of boron, iron, and hydrogen peroxide on material metabolism in plants: *Akad. Nauk SSSR, Doklady* v. 94, p. 157-160.
- Shoup, C. S., and Boykin, J. T., 1931, The insensitivity of *Paramecium* to cyanide and effects of iron on respiration: *Jour. Gen. Physiology*, v. 15, p. 107-118.
- Siddappa, G. S., and Subrahmanyam, V., 1935, Role of organic matter in plant nutrition, part 9—Oxidation of organic matter in the soil and plant assimilation: *Indian Acad. Sci. Proc.*, 1B, p. 928-937.
- Sideris, C. P., and Young, H. Y., 1956, Pineapple chlorosis in relation to iron and nitrogen: *Plant Physiology*, v. 31, p. 211-222.
- Sidorin, M. I., 1914, On the assimilation of iron by plants: *Izv. Rezul't Veget. Opytov Lab. Rabot (Rec. Trav. Lab. Agron.) Moskovskii Sel'Skokhoziai-stvennyi Institut*, v. 10, p. 241-257.
- Silverman, L., and Valenzuela, C., 1946, Filter paper collection of cadmium and iron fumes in air: *Jour. Indus. Hygiene Toxicology*, v. 28, p. 107-111.
- Simakov, V. N., and Isakova, A. I., 1933, The influence of saturating the absorbing complex with iron on the biochemical processes in the soil: *Pochvovedenie*, v. 28, p. 170-184.
- Simons, L. H., Monaghan, P. H., and Taggart, M. S., Jr., 1953, Aluminum and iron in Atlantic and Gulf of Mexico surface: *Anal. Chemistry*, v. 25, p. 989-990.
- Singh, R. M., 1955, Effect of iron on leaf area of wheat plant: *Sci. and Culture [India]*, v. 21, p. 318-319.
- 1956, Effect of iron on leaf growth and chlorophyll content of wheat plants: *Sci. and Culture [India]*, v. 22, p. 229-231.
- Sisakyan, N. M., and Melik-Sarkisyan, S. S., 1956, The nature of chloroplast proteins: *Biokhimiya*, v. 21, p. 329-339.
- Skadovskii, S. N., Messineva, M. A., and Uspenskaya, V. I., 1955, Studies on microorganisms causing the appearance and disappearance of odors and tastes in water: *Vsesoyuznogo Gidrobiol. Obshchestva Trudy*, v. 6, p. 23-37.
- Skeen, J. R., 1930, Experiments with *Trianea* on antagonism and absorption: *Plant Physiology*, v. 5, p. 105-118.

- Skopintsev, B. A., and Krylova, L. P., 1955, Dynamics of organic substances in natural waters: *Vsesoyuznogo Gidrobiol. Obshchestva Trudy*, v. 6, p. 38-45.
- Slobodchikov, B. Y., and Stroikina, V. G., 1953, Effect of nitrogen, phosphorus, and iron on the development of phytoplankton in Sevan Lake: *Akad. Nauk Armyanskoi SSR Izv., Biol. Sel'skokhozyaistvennye Nauki*, v. 6, no. 7, p. 3-18.
- Smirnov, A. A., 1955, Formation of underground carbonic acid waters: *Gidrokhimicheskie Materialy*, v. 24, p. 101-107.
- Smith, G. F., McCurdy, W. H., Jr., and Diehl, Harvey, 1952, Colorimetric determination of iron in raw and treated municipal water supplies by use of 4,7-diphenyl-1, 10-phenanthroline: *Analyst*, v. 77, p. 418-422.
- Smith, G. F., and Richter, F. P., 1944, Substituted 1,10-phenanthroline ferrous complex oxidation-reduction indicators. Potential determinations as a function of acid concentration: *Indus. Eng. Chemistry, Anal. Ed.*, v. 16, p. 580-581.
- Smith, G. M., 1950, *Fresh-water algae of the United States*: 2d ed., New York, McGraw-Hill, p. 16.
- Smith, H. J., McIlrath, W. J., and Bogorad, L., 1957, Effects of iron deficiency on flowering of *Xanthium*: *Bot. Gaz.*, v. 118, p. 174-179.
- Solovev, A. V., 1940, The formation of acid and corrosion of metals in certain soils: *Zhur. Khim. Promyshlennosti*, v. 17, no. 9, p. 42-45.
- Somers, I. I., and Shive, J. W., 1942, The Fe-Mn relation in plant metabolism: *Plant Physiology*, v. 17, p. 582-602.
- Sorokin, Y. I., 1955, Determination of the magnitude of chemosynthesis in the waters of Rybinsk reservoir by means of carbon-14: *Akad. Nauk SSSR, Doklady*, v. 105, p. 1343-1345.
- Sorokina, V. A., 1938, Exchange of substance between slime and water as influenced by the formation of a bacterial film on the surface of the slime: *Limnological Sta. Kosin Microbiol. Lab., Moscow State Univ.; Microbiology*, v. 7, p. 579-591 [English summary].
- Speirs, M., 1954, Effect of environment on the iron content of plants: *Southern Coop. Ser. Bull* 36, p. 155-175.
- Speller, F. N., 1922, Control of corrosion of deactivation of water: *Jour. Franklin Inst.*, v. 193, p. 523.
- 1951, *Corrosion, Causes and prevention*: 3d ed., New York, McGraw-Hill, 686 p.
- Spizizen, J., 1943, Biochemical studies of the phenomenon of virus reproduction, part 2—Studies on the influence of compounds and metabolic significance on the multiplication of bacteriophage: *Jour. Infectious Diseases*, v. 73, p. 222-228.
- Starkey, R. L., 1945, Transformations of iron by bacteria in water: *Am. Water Works Assoc. Jour.*, v. 37, p. 963-984.
- Starkey, R. L., and Wight, K. M., 1943, Soil areas corrosive to metallic iron through activity of anaerobic sulfate-reducing bacteria: *Am. Gas Assoc. Monthly*, v. 25, no. 5, p. 223-228.
- Steel, E. W., and Gloyne, E. F., 1954, Oxidation ponds—radioactivity uptake and algae concentration: *U.S. Atomic Energy Comm. no. 275*, p. 10-37.
- 1955, Concentration of radioactivity in oxidation ponds: *Sewage and Indus. Wastes*, v. 27, p. 941-956.

- Steele, C. C., 1949, An introduction to plant biochemistry: 2d ed., London, G. Bell and Sons, 346 p.
- Steigmann, A., 1946, Metals in gelatin: Food, v. 15, p. 105-106.
- Steinberg, R. A., 1919, A study of some factors in the chemical stimulation of the growth of *Aspergillus niger*: Am. Jour. Botany, v. 6, p. 330-372.
- 1920, Effect of zinc and iron compared with that of uranium and cobalt on growth of *Aspergillus*: Bot. Gaz., v. 70, p. 465-468.
- 1935a, The nutritional requirements of the fungus *Aspergillus niger*: Bull. Torrey Bot. Club, v. 62, p. 81-90.
- 1935b, Nutrient solution purification for removal of heavy metals in deficiency investigations with *Aspergillus niger*: U.S. Dept. Agriculture Jour. Agr. Research, v. 51, p. 413-424.
- 1936, Some effects of the heavy metals essential for the nutrition of *Aspergillus niger* upon its growth: Am. Jour. Botany, v. 23, p. 227-231.
- 1941, Use of *Lemna* for nutrition studies on green plants: U.S. Dept. Agriculture Jour. Agr. Research, v. 62, p. 423-430.
- 1943, Use of *Lemna* as test organism: Chronica Botanica, v. 7, p. 420.
- 1946, Mineral requirement of *Lemna minor*: Plant Physiology, v. 21, p. 42-48.
- 1950, Growth on synthetic nutrient solutions of some fungi pathogenic to tobacco: Am. Jour. Botany, v. 37, p. 711-714.
- Steinkamp, R., Dubach, R., and Moore, C. V., 1955, Iron transportation and metabolism, part 8—Absorption of radioiron from iron-enriched bread: Archives Int. Medicine, v. 95, p. 181-193.
- Stern, P., Kosak, R., Misirlija, A., and Hukovic, S., 1955, Influence of the intestinal flora on iron absorption: Naunyn-Schmiedeberg's Archiv Expt. Pathologie Pharmacologie, v. 225, p. 162-163.
- Steward, F. C., 1948, Metabolism and salt uptake by plants: Brookhaven Conf. Rept. BNL-C-4, p. 94-103.
- Stewart, I., and Leonard, C. D., 1956, Studies of chelated plant nutrients with radioactive isotopes: U.S. Atomic Energy Comm. TID-7512, p. 245-251.
- Stobe, G., 1956, Oxidation-reduction potentials of soils: Latvijas SSR Zinatnu Akad. Vestis 6, p. 41-47.
- Stokes, J. L., 1954, Studies on the filamentous sheathed iron bacterium *Sphaerotilus natans*: Jour. Bacteriology, v. 67, p. 278-291.
- Stoklasa, J., 1898, The physiological function of iron in plants: Acad. sci. [Paris] Comptes rendus, v. 127, no. 5, p. 282-283.
- Stoltenberg, H. A., 1949, Oxygen concentration effect on flood water iron pickup: World Oil, v. 128, no. 9, p. 154, 156.
- Strakhov, N. M., 1948, Distribution of iron in lake sediments and seawater and the factors of its control: Akad. Nauk SSSR Izv., Ser. geol., no. 4, p. 3-50.
- Stumm, W., 1956, Calcium carbonate deposition at iron surfaces: Am. Water Works Assoc. Jour., v. 48, p. 300-310.
- Sugawara, K., 1934, Liesegang's stratification developed in the diatomaceous gythia from Lake Haruna, and problems related to it: Chem. Soc. Japan Bull., v. 9, p. 402-409.
- Supniewski, J., 1951, The role of organically combined iron in animal organisms: Farmacia Polska, v. 7, p. 197-202.
- Susplugas, J., Pelinard, J., and Danoy, M., 1951, Mineral content and plant development: Soc. Pharmacie Montpellier Travaux v. 2, p. 25-33.
- Svendenius, G., 1929, Gravimetric determination of iron in organic substances: Acta Paediatrica, v. 9, p. 1-8.

- Taboury, F. J., and Mangin, C., 1948, Electrolytic dissociation of molecular addition complexes at the hydrogen (bonding) bridge: Soc. Chim. France Bull., p. 47-52.
- Tanaka, M., 1953, Chemical study of the mineral metabolism of lakes: Nagoya Univ., Jour. Earth Sci., v. 1, p. 119-134.
- 1954, The nature of iron contained in lake waters: Chem. Soc. Japan Bull., v. 28, p. 89-93.
- Temple, K. L., and Colmer, A. R., 1951, The autotrophic oxidation of iron by a new bacterium: *Thiobacillus ferrooxidans*: Jour. Bacteriology, v. 62, p. 605-611.
- Thiele, H., 1928, Iron and manganese in water: Gas- u. Wasserfach, v. 71, p. 289-290.
- Thomas, G. and Brossard, J., 1956, Reaction between iron salts and tannins from the chestnut (*Castanea sativa*): Chemistry and Industry [London], p. 440-441.
- Thompson, M. K., 1940, The reaction between iron and water in the absence of oxygen: Electrochem. Soc. Trans., v. 78, 5 p.
- Thompson, T. G., and Bremner, R. W., 1935a, The determination of iron in seawater: Conseil Permanent Internat. Explor. Mer., Jour. conseil, v. 10, p. 33-38.
- 1935b, The occurrence of iron in the water of the northeast Pacific Ocean: Conseil Permanent Internat. Explor. Mer., Jour. conseil, v. 10, p. 39-47.
- Thompson, T. G., Bremner, R. W., and Jamieson, I. M., 1932, Occurrence and determination of iron in seawater: Indus. Eng. Chemistry, Anal. Ed., v. 4, p. 288-290.
- Thorne, D. W., and Walker, R. H., 1936, Physiological studies on *Rhizobium*, part 6—Accessory factors: Soil Sci., v. 42, p. 231-240.
- Thornton, H. G., 1956, Ecology of microorganisms in soil: Royal Soc. [London] Proc., v. B145, p. 364-374.
- Tillmans, J., and Klarmann, B., 1924, The solution of iron by oxygen-free natural waters: Zeitschr. Angew. Chemie, v. 36, p. 94-97, 103-104, 113-115.
- Todt, F., 1934, The increase of rate of solution of iron by oxidizing and reducing substances: Zeitschr. Elektrochemie, v. 40, p. 536-541.
- 1950, The reaction of atmospheric oxygen dissolved in water on metal surfaces: Chemiker-Zeitung, v. 74, p. 254-256.
- Toth, S. J., 1937, Anion adsorption of soil colloids in relation to changes in free iron oxides: Soil Sci., v. 44, p. 299-314.
- Trojanowski, J., 1954, The biological activity of some humic fractions: Soc. Botanicorum Poloniae Acta, v. 23, p. 143-160.
- Troug, E., 1946, Soil reaction influence on availability of plant nutrients: Soil Sci. Soc. America Proc., v. 11, p. 305-308.
- Tsujimura, K., and Ikeda, F., 1957, Iron ethylenediaminetetraacetate chelate for water culture: Nippon Dojo- Hiriyogaku Zasshi, v. 27, p. 479-480.
- Tullis, E. C., and Cralley, E. M., 1936, Chlorosis of rice induced by iron deficiency: Phytopathology, v. 26, p. 111.
- Twyman, E. S., 1951, The iron and manganese requirements of plants: New Phytologist, v. 50, p. 210-226.
- Tysarowski, W., 1955, The complex ion from iron and ethylenediaminetetraacetic acid as an electron-transporting catalyst in biological model experiments: Acad. Polonaise Sci. Bull., Cl. 2, v. 3, p. 201-205.

- Ueda, H., 1956, Effects of iron on the growth of lowland rice (*Oryza sativa*): Tottori Univ. Jour. Fac. Agriculture, v. 2, p. 117-142.
- Ueda, K., Yannaoka, M., and Sakichira, N., 1956, Aged paddy soils, part 3—Effect of material high in iron oxide on rice-plant growth in aged paddy soils: Jour. Sci. Soil Manure, Japan, v. 27, p. 69-71.
- Umemoto, S., and Mifune, M., 1953, Chemical studies on thermophilic algae, part 2—Inorganic constituents: Okayama Univ. Repts. Balneological Lab., v. 12, p. 11-14.
- Unti, O., 1943, Fe content of waters carrying *Anopheles* or *Anopheles* without malaria in the valley of Paraiba: Arquivos Higiene Saude Publica [Sao Paulo], v. 8, no. 19, p. 105-117.
- Upadhyaya, M. P., 1955, Calcium content of fishery waters of Uttar Pradesh: Indian Jour. Veterinary Sci., v. 25, p. 341-345.
- Urbach, C., 1934, Step-photometric microanalysis of drinking and service water, part 12—Iron determination: Mikrochemie, v. 15, p. 207-226.
- Urbain, O. M., and Miller, J. N., 1930, The effect of iron on the accuracy of the determination of dissolved oxygen by the Winkler method: Am. Water Works Assoc. Jour., v. 22, p. 1261-1265.
- Urs, M. K., Natarajan, C. P., and Bhatia, D. S., 1955, Mineral nutrients in coffee brew: Jour. Sci. Indus. Research [India], v. 14c, p. 60.
- Uspenski, E. E., 1927, Eisen als Faktor fur die Verbreitungniederer Wasserpflanzen: Pflanzforschung, 104 p.
- Valyashko, M. G., 1955, Principal chemical types of water and their formation: Akad. Nauk SSSR Doklady, v. 102, p. 315-318.
- Van Beneden, G., 1947, Iron in water from gravel: Technique l'eau [Brussels], p. 11-12.
- 1948, Mineral waters: Technique l'eau [Brussels], v. 2, no. 4, p. 23-24.
- 1956a, Determination of "humic iron;" Application of this method to the peaty waters and the creeks of the Ardennes: Centre Belge etude doc. eaux [Liège] Bull. 31, p. 12-14.
- 1956b, The behavior of humic substances in water. Humic acid flocculation in peat-bog waters, part 2—Action of electrolytes: Centre Belge étude doc. eaux [Liège] Bull. 33, p. 173-181.
- Van de Spek, J., 1948, Iron in soil: Chem. Weekblad, v. 44, p. 493-499; 505-511.
- Vanstone, F. H., 1951, Quantitative method of flame spectrographic analysis: Ann. Rept. East Malling Research Sta., Kent, p. 122-125.
- Varkov, S. P., Kulakov, E. V., and Kaurichev, I. S., 1950, The formation of ferrous iron and the peculiarities of the phosphate regime in sod-podzolized soils: Pochvovedenie p. 466-475.
- Vialard-Goudon, A., and Richard, C., 1955, Reduction by iron, of NO_3 ions to NO_2^- and NH_4^+ ions in acidic tropical waters: Acad. sci [Paris] Comptes rendus, v. 241, p. 978-980.
- Vidal, J. L., 1938, Treatment of calcareous chlorosis of fruit trees: Acad. agriculture, France, Comptes rendus, v. 24, p. 160-163.
- Villeret, S., 1953, A hydrobiological study of ponds in the vicinity of Rennes, part 1—Mare des Buttes: Soc. sci. Bretagne Bull., v. 28, p. 97-111.
- 1954a, The biology of algae in *Sphagnum* peatbogs: Soc. sci. Bretagne Bull., v. 29, spec. no., p. 1-246.
- 1954b, The algal vegetation of the Peat bog of Nesnay en Plouneour [Finistere] Soc. sci. Bretagne Bull., v. 29, p. 17-31.
- 1954c, The biology of algae in *Sphagnum* peatbogs: Soc. sci. Bretagne Bull., v. 29, p. 32.

- Villeret, S., 1954d, The presence of two supposedly rare fresh-water algae, *Cycloneis annularis* and *Glocotaenium loitlesbergerianum*, in Brittany: Soc. sci. Bretagne Bull., v. 29, p. 33-40.
- 1954e, The use of "complexous" in limnology: Soc. sci. Bretagne Bull., v. 29, p. 41-47.
- Vinberg, G. G., 1955, Significance of photosynthesis in enrichment of water with oxygen in self-purification of contaminated waters: Vsesoyuznogo Hidrobiol. Obshchestva Trudy, v. 6, p. 46-69.
- Virtanen, A. I., Jorma, J., Linkola, H., and Linnasahmi, A., 1947, The relation between nitrogen fixation and leghemoglobin content of leguminous root nodules: Acta Chemica Scandinavica, v. 1, p. 90-111.
- Voichevich, P. G., 1940, Control of weeds by iron sulfate: Sovetskaya Agronomiya, no. 5, p. 79-80.
- Volkova, O. Y., 1939, Ferrobacteria of caucasian mineral springs and their role in the formation of ferrous deposits: Microbiology [U.S.S.R.], v. 8, p. 863-886.
- Voronkov, P. P., 1953, Basic features of biogenous compounds in the reservoirs of the Volga system, in connection with their primary production: [U.S.S.R.] Gosudarst. Gidrol. Inst., no. 37, p. 62-80.
- Vouk, L., 1936, The iron stores of blue algae: Mikrochemie, p. 439-446.
- Voznesenskii, S. A., and Nagatkin, I. T., 1934, The velocity of the reaction of iron with cupric sulfate: Zhur, priklad khimii v. 7, p. 51-50.
- Walker, J. B., 1954, Inorganic micronutrient requirements of *Chlorella*, part 2—Quantitative requirements for iron, manganese, and zinc: Archives Biochemistry and Biophysics, v. 53, p. 1-8.
- 1956, Strontium inhibition of calcium utilization by a green alga: Archives Biochemistry and Biophysics, v. 60, p. 264-265.
- Walker, W. H., Cederholm, A. M., and Bent, L. N., 1907, The corrosion of iron and steel: Am. Chem. Soc. Jour., v. 29, p. 1251-1264.
- 1908, The corrosion of iron and steel: Am. Chem. Soc. Jour., v. 30, p. 473-474.
- Wallace, A., and Ashcroft, R. T., 1956, Ammonium lignin sulfonate as a chelating agent for supplying soluble iron to plants: Soil Sci., v. 82, p. 233-236.
- Wallace, A., and Lunt, O. R., 1956, Reactions of some iron, zinc, and manganese chelates in various soils: Soil Sci. Soc. America Proc., v. 20, p. 479-482.
- Wallace, A., North, C. P., Mueller, R. T., Shannon, L. M., and Hemaidan, N., 1955, The behavior of chelating agents in plants: Am. Soc. Hort. Sci. Proc., v. 65, p. 9-16.
- Wallace, T., 1929, Investigations on chlorosis of fruit trees, part 4—The control of lime-induced chlorosis in the field: Pomology and Hort. Sci. Jour., v. 7, no. 4, p. 251-269.
- 1954, Effects of mineral deficiencies on crop plants: Cong. Internat. Botan., 8th, Paris, Rap. et commun., v. 8, sect. 11-12, p. 87-91.
- Wallbach, G., 1932, Microscopically visible endogenous iron metabolism: Klinische Wochenschr., v. 11, p. 353-356.
- Wann, F. B., 1930, Chlorosis yellowing of plants: Cause and Control: Utah State Agr. Coll. Agr. Expt. Sta. Circ. 85, 12 p.
- Warburg, O., 1925, Character of respiration, its explanation: Pharmazeut. Monatsh., v. 6, p. 105-106.
- Warburg, O., Negelein, E., and Haas, E., 1933, Spectroscopic demonstration of the oxygen-transferring enzyme in the presence of cytochrome: Biochem. Zeitschr., v. 266, p. 1-8.

- Ward, H. B., and Whipple, G. C., 1918, *Fresh-water biology*: 1st ed., New York, Wiley and Sons, Inc., 1111 p.
- Waring, W. S., and Werkman, C. H., 1942, Growth of bacteria in an iron-free medium: *Archives Biochemistry*, v. 1, p. 303-310.
- 1943, Iron requirements of heterotrophic bacteria: *Archives Biochemistry*, v. 1, p. 425-433.
- Waser, E., Husmann, W., and Blochliger, G., 1937, *The limmat: Vom Wasser*, v. 12, p. 181-245.
- Wasser, O., 1938, Iron in water and the iron bacteria: *L'Eau* v. 31, p. 103-105.
- Watkins, J. W., 1954, Analytical methods of testing waters to be injected into subsurface oil-productive strata: U.S. Bur. Mines, Rept. Inv. 5031, 29 p.
- Watson, G. H., and Bollen, W. B., 1952, Effect of copper sulfate weed treatment on bacteria in lake bottoms: *Ecology*, v. 33, no. 4, p. 522-529.
- Weart, J. G., and Margrave, G. E., 1957, Oxidation-reduction measurements applied to iron removal: *Am. Water Works Assoc. Jour.*, v. 49, p. 1223-1236.
- Weinstein, L. H., Robbins, W. R., and Perkins, H. F., 1954, Chelating agents and plant nutrition: *Science*, v. 120, p. 41-43.
- Werescagin, G. J. Anickova, N. J., and Forsch, T. B., 1931, Methods of hydrochemical analysis in limnology: *Achiv Hydrobiologie*, v. 23, p. 1-64; 167-230.
- Weston, R. S., 1936, Some problems of deferrization and demanganization: *New England Water Works Assoc. Jour.*, v. 50, p. 231-238.
- Whatley, F. R., Ordin, L., and Arnon, D. I., 1951, Distribution of micronutrient metals in leaves and chloroplast fragments: *Plant Physiology*, v. 26, p. 414-418.
- Wheatly, W., 1946, High iron content of water of Crystal City, Missouri: *Public Works*, v. 77, no. 1, p. 36-38.
- Whipple, G. C., Fair, G. M., and Whipple, M. C., 1927, *The microscopy of drinking water*: New York, Wiley and Sons, 586 p.
- White, H. E., 1937, Study of the effect of plant nutrients, soil reaction, and light on gardenias: *Massachusetts Agr. Expt. Sta. Ann. Rept.*, Bull. 339, p. 64-65.
- Wickert, K., and Pilz, H., 1950-51, Trace analysis in water from power stations: *Vom Wasser*, v. 18, p. 303-309.
- Wiebosch, W. A., 1947, Chlorosis in North Holland: Its identification and control: [Netherlands] *Direct. Tuinbouwvoorlichting Mededeel.* v. 10, p. 556-565.
- Willcox, L. V., 1948, The quality of water for irrigation use: U.S. Dept. Agriculture Tech. Bull. 962, 40 p.
- Will, J. J., and Vilter, R. W., 1954, The absorption and utilization of an iron chelate in iron deficient patients: *Jour. Lab. Clinical Medicine*, v. 44, p. 499-505.
- Willcomb, G. E., 1936, Iron and manganese in water: *Am. Water Works Assoc. Jour.*, v. 28, p. 1896-1909.
- Williams, R. J. P., 1953, Metal ions in biological systems: *Cambridge Philos. Soc. Biol. Rev.*, v. 28, p. 381-415.
- Williams, R. P., Green, J. A., and Rappoport, D. A., 1956, Evidence for the incorporation of iron and calcium into the pigments of *Serratia marcescens*: *Science*, v. 123, p. 1176-1177.
- Willis, L. G., 1936, Evidences of the significance of oxidation-reduction equilibrium in soil fertility problems: *Soil Sci. Soc. America Proc.*, v. 1, p. 291-297.
- Willis, L. G., and Carrero, J. O., 1921, Chlorosis of rice: *Puerto Rico Univ. Agr. Expt. Sta. Bull.*, p. 8-9.

- Willis, L. G., and Piland, J. R., 1936, The function of copper in soils and its relation to the availability of iron and manganese: U.S. Dept. Agriculture Jour. Agr. Research, v. 52, p. 467-476.
- 1937, Some recent observations on the use of minor elements in North Carolina agriculture: Soil Sci., v. 44, p. 251-263.
- Wind, J., 1951, Traces of elements in agriculture: Maandblad Landbouwworlichtingsdienst, v. 8, p. 223-233.
- Winogradsky, S., 1922, Iron bacteria as inorgoxidants: Centralbl. Bakteriologie Parasitenkunde, Abt. 2, v. 57, p. 1-21.
- Winsor, H. W., 1939, Mineral content of vegetable crops with special reference to iron: Florida Agr. Expt. Sta. Ann. Rept., p. 85-86.
- Winter, O. B., 1936, Report on (the analysis of) plants: Assoc. Official Agr. Chemists Jour., v. 19, p. 359-365.
- Winters, E., 1940, The migration of iron and manganese in colloidal systems: Illinois Agr. Expt. Sta. Bull. 472, p. 101-143.
- Wolken, J. J., and Mellon, A. D., 1956, The relation between chlorophyll and the cortenoids in the algal flagellate, *Euglena*: Jour. Gen. Physiology, v. 39, p. 675-685.
- Woodcock, A. H., 1955, Bursting bubbles and air pollution: Sewage and Indus. Wastes, v. 27, p. 1189-1192.
- Wort, D. J., 1955, The seasonal variation in chemical composition of *Macrocystis integrifolia* and *Nereocystis luekeana* in British Columbia coastal waters: Canadian Jour. Botany, v. 33, p. 323-340.
- Yamagata, K., Oda, M., and Ando, T., 1955, Variation of *Aspergillus oryzae*, part 7—Variation induced by chemicals: Jour. Fermentation Technology [Japan], v. 33, p. 369-373.
- Yamasaki, T., 1952, Excess moisture injury of upland crops in overmoist soil from the viewpoint of soil chemistry and plant physiology: Natl. Inst. Agr. Sci. Bull., ser. B, no. 1, 92 p.
- Yarilova, E. A., 1947, The role of lithophilic lichens in weathering crystalline bedrock: Pochvovedenie, p. 533-548.
- Yoshii, G., and Watabe, A., 1956, A method of removing fission products of atomic nucleus by means of plankton: Kagaku [Tokyo], v. 26, p. 363-364.
- Yoshimura, F., 1950, Physiological studies in lemnaeous plants, parts 3 and 4: Bot. Mag. Tokyo, v. 63, p. 63-69.
- Yoshimura, S., 1931a, Stratification of iron and manganese in lake water in Japan: Japanese Jour. Geology Geography, v. 9, nos. 1-2, p. 61-69.
- 1931b, The hydrogen-ion concentration of the lake water in Japan: Imp Acad. Tokyo Proc., v. 7, p. 195-197.
- 1931c, Seasonal variation of iron and manganese in the water of Taka suka-numa, Saitama: Japanese Jour. Geology Geography, v. 8, p. 269-279.
- 1934, Anohaline stratification of the chemical constituents of Lake Osore-sanko, Aomori Prefecture, Japan: Imp. Acad. Tokyo Proc., v. 10, p. 475-478.
- 1936, Contributions to the knowledge of iron dissolved in the lake waters of Japan, part 2: Japanese Jour. Geology Geography, v. 13, p. 39-56.
- Yust, W., 1951, Iron bacteria: Chicago, Ill., Encyclopedia Britannica, Inc., v. 2, p. 908.
- Zapffe, C., 1931, Deposition of manganese: Econ. Geology, v. 26, p. 799-832.
- 1933, Catalysis and its bearing on origin of Lake Superior iron-bearing formation: Econ. Geology, v. 28, p. 751-772.

- Zhivopistsev, V. P., and Minina, V. S., 1954, Colorimetric determination of iron in natural waters with 2-furyldiantipyrinylmethane: Molotovskogo Gosudarstvennogo Univ. Imeni A. M. Gor'kogo, Uchenye Zapiski, v. 8, no. 3, Mat. Fiz. Khim., p. 37-41.
- Zicker, E. L., Berger, K. C., and Hasler, A. D., 1956, Phosphorus release from bog lake muds: Limnology Oceanography, v. 1, no. 4, p. 296-303.
- Zonn, S. V., 1945, Chemical composition of ground waters as dependent on soil formation: Acad. sci. U.R.S.S. Comptes rendus, v. 48, p. 197-199.

INDEX

	Page		Page
<i>Acromatium oxaliferum</i>	132	<i>Crenothrix fusca</i>	131, 132
Actinomycetes.....	113, 129	<i>polyspora</i>	132, 133
<i>Aerobacter aerogenes</i>	124	sp.....	131, 132, 133, 140
<i>indologenes</i>	124	Desert varnish.....	134
Agar-agar, used in experiments.....	115; fig. 12	Desmidiaceae.....	126, 140
Amines.....	138	Diatoms.....	126, 140
<i>Anabaena lemmermanni</i>	126	<i>Elodea</i>	136
<i>Anthophysa vegetans</i>	131, 133	<i>Enteromorpha</i>	137
<i>Aphanizomenon flos-aquae</i>	126	<i>Escherichia coli</i>	125
<i>Asellus aquaticus</i>	133	<i>Euglena</i> sp.....	126
<i>Aspergillus niger</i>	124	<i>Ferrobacillus ferrooxidans</i>	125
<i>oryzae</i>	124	<i>Flavobacterium halobium</i>	125
<i>Asterionella formosa</i>	126	<i>Fucus vesiculosus</i>	140
<i>japonica</i>	126	<i>Fusarium lini</i>	124
<i>Azotobacter chroococcum</i>	124	<i>moniliforme</i>	124
sp.....	124	<i>orthoceras</i>	124
<i>vinelandii</i>	124	<i>oxysporum</i>	124
Bacilliform bacterium.....	125	<i>poae</i>	124
<i>Bacillus bruntzii</i>	124	<i>scirpi</i>	124
<i>megatherium</i>	123	sp.....	124
<i>prodigiosus</i>	124	<i>udum</i>	124
Bacteria, effect on iron content.....	114, 123, 138, 142	<i>vasinfectum</i>	124
hemophilic.....	125	<i>Gallionella ferruginea</i>	132
iron-using.....	124-125, 129-134	Gametophyte.....	113, 134
marine.....	125, 132	<i>Helminthosporium sativum</i>	124
produce blackbeet disease.....	125	Human consumption of water, domestic.....	112, 118, 128
soap-discoloring.....	125	effect of iron content.....	112
<i>Bacterium precipitatum</i>	131, 132	industrial.....	112
Benthos.....	113	Hydrophytes, definition.....	113
<i>Bikosoeca petiolata</i>	131	effect on iron content.....	114, 118, 134-141
Biological terms.....	113	identification.....	112
<i>Brucella abortus</i>	124	<i>Iridaea</i> sp.....	140
Cambium.....	113, 138	Iron bacteria, calcium-metabolizing.....	132
<i>Cercospora nicotinae</i>	124	classification.....	129
<i>Chaetoceras curvisetius</i>	126	harmful effects.....	131
Chelating agents.....	115, 119, 134	manganese-metabolizing.....	131
<i>Chilomonas paramecium</i>	125	methods of control.....	133
<i>Chlamydomonas</i> sp.....	126	organic iron metabolizing.....	132
<i>Chlamydothrix ochracea</i>	129	sulfur-metabolizing.....	132
<i>sideropons</i>	129	Iron concentrations, effect on plants.....	117, 118
<i>Chlorella pyrenoidosa</i>	126, 141	Iron content of natural water, factors affect- ing.....	112, 144
sp.....	126, 140	Iron cycle.....	111, 112, 114
<i>vulgaris</i>	126	Iron solubility, affected by organic matter.....	119-121
<i>Chlorobium limicola</i>	125, 132	Irrigation.....	143
<i>Chlorobium thiosulfatophilum</i>	125, 132	<i>Klebsiella pneumoniae</i>	125
Chloroplast.....	113	<i>Laminaria digitala</i>	140
Chromosome.....	113	<i>Lemna major</i>	127
<i>Cladophora glomerata</i>	126	<i>minor</i>	127
sp.....	140, 141		
<i>Cladothrix dichotoma</i>	129, 132		
<i>Clonothrix fusca</i>	129		
<i>Clostridium tetani</i>	125		
<i>Coelospaerium</i> sp.....	126		
Coenocyte.....	113		

<i>Lemna</i> —Continued	Page		Page
<i>polyrhiza</i>	127	<i>Potamogeton foliosus</i>	127
sp.....	127	sp.....	127, 140
<i>Leptothrix echinata</i>	132	Protozoa.....	113, 125
ochracea.....	129, 132, 133	<i>Pseudomonas aeruginosa</i>	125
sp.....	132, 141	<i>cattleyae</i>	125
<i>trichogenes</i>	132	<i>cutirubra</i>	125
Littoral.....	113	<i>ferrugineum</i>	132
<i>Lyngbia confervoides</i>	126	<i>salinaria</i>	125
<i>mainsecula</i>	126	Pyrite, oxidation.....	119
<i>Macrocystis integrifolia</i>	126	<i>Pythium irregulare</i>	124
Mercaptans.....	138	Reduction of iron.....	121, 122, 123
Meristem.....	113, 138	Respiration, aerobic.....	118, 119
Mesophyll.....	113, 138	anaerobic.....	118, 119, 135
<i>Microcystis aeruginosa</i>	126, 140	effect of iron concentration.....	117, 118
Mycelium.....	113	<i>Rhizobium meliloti</i>	125
See also Plants, fungi.		sp.....	125
<i>Mycobacterium phlei</i>	125	<i>trifolii</i>	125
<i>tuberculosis</i>	125	<i>Rhizoctonia solani</i>	124
Natural water, iron content.....	111	<i>Rhodopseudomonas</i> sp.....	125
<i>Nereocystisluetkeana</i>	126	<i>Rhodymenia</i> sp.....	140
<i>Neurospora crassa</i>	124	<i>Sarcina littoralis</i>	125
<i>Nitzschia closterium</i>	126	<i>Sargassum siliquastrum</i>	140
<i>Nostoc flagelliforme</i>	140	<i>Sclerotium rolfsii</i>	124
<i>Oedogonium</i> sp.....	140	<i>Serratia marcescens</i>	125
<i>Oryza sativa</i>	127	<i>Siderocapsa</i> sp.....	132
<i>Oscillatoria agarhdia</i>	126	<i>Siderodendron manganiferum</i>	132
<i>rubescens</i>	126	<i>Siphomonas fritschii</i>	132
Oxidation of iron.....	114, 119-122	<i>Sphaerotilus natans</i>	129, 131, 133
<i>Paramecium caudatum</i>	125	<i>Spirophyllum ferrugineum</i>	132
Pelagic.....	113	sp.....	132
Phage.....	113, 125	Sporophyte.....	114, 134; fig. 14
Phloem.....	113, 135, 138	<i>Streptomyces griseus</i>	124
<i>Phormidium ambiguaum</i>	126	Tannins, occurrence.....	137, 138, 144
Photosynthesis, empirical formula.....	118	<i>Thielaviopsis basicola</i>	124
iron-concentration effects.....	117, 118, 137	<i>Thiobacillus denitrificans</i>	125
Plankton.....	113	<i>ferrooxidans</i>	125, 132
Plants, algae.....	113, 114, 115, 118	<i>thiooxidans</i>	125
algae attached.....	114, 115, 137	<i>Toxothrix</i> sp.....	132
blue-green.....	126	<i>Trachelomonas</i> sp.....	133
brown.....	126, 140	<i>Trichophyton mentagrophytes</i>	124
green.....	126	<i>rubrum</i>	124
iron-using.....	126, 134, 137, 141	<i>Trypanosoma</i> sp.....	125
unattached.....	115, 137	<i>Typha latifolia</i>	127
aquatic, characteristics.....	134-136	<i>Ulothrix</i> sp.....	126
crop.....	127	<i>zonata</i>	141
dictyyledon.....	113; fig. 17	Use of iron, by animals.....	111, 117, 118, 125, 128, 138
ferns.....	114	by plants.....	111, 115, 117,
flowering.....	114, 127	118, 124, 125, 126, 127, 128, 134, 138, 143	
iron-using.....	127, 137, 138	<i>Ustilago sphaerogena</i>	124
fungi.....	113, 114, 124, 129, 133	<i>Vibrio desulphuricans</i>	132
molds.....	124	Virus.....	113, 114, 125
lichens.....	113, 115, 134	Water, analysis.....	144, 145-149
monocotyledon.....	113, 140	Water color, in relation to organic content.....	119,
mosses.....	113, 115	121, 131	
nonvascular.....	113	Xylem.....	114, 135
soil-rooted.....	112, 114, 127	Yeasts.....	125
vascular.....	114	<i>Zygnemaceae</i>	140
water-rooted.....	112, 114, 127		
See also Hydrophytes.			
<i>Pleurobrachia</i> sp.....	125		